

MAP2K1 recombinant antibody

MAP2K1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243994	Rabbit	Human, Rat	Immunoprecipitation, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

MAP2K1; Monoclonal Recombinant Antibody; MAP2K1 Antibody; Dual specificity mitogen-activated protein kinase kinase 1 (MAP kinase kinase 1) (MAPKK 1) (MKK1) (EC 2.7.12.2) (ERK activator kinase 1) (MAPK/ERK kinase 1) (MEK 1); MEK1 PRKMK1; MAP2K1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	9B6
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for MAP2K1 recombinant antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human MEK1
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IP (Immunoprecipitation)

Immunoprecipitating MAP2K1 in HeLa whole cell lysate Lane 1: Rabbit control IgG instead of in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000) Lane 2: HeLa whole cell lysate 500ug Lane 3: HeLa whole cell lysate (10ug)



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human glioma cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)
 Western BlotPositive WB detected in: HeLa whole cell lysate, 293 whole cell lysate, MCF-7 whole cell lysate, 293T whole cell lysate, A549 whole cell lysate, U251 whole cell lysate, Rat brain tissueAll lanes: MAP2K1 antibody at 1:2000SecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 44, 41 kDaObserved band size: 44 kDa

ADDITIONAL INFORMATION

Related Product Information for MAP2K1 recombinant antibody	Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.
Product Categories/Family for MAP2K1 recombinant antibody	Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	5579478
NCBI GeneID	5604
NCBI Accession #	NP_002746.1
NCBI GenBank Nucleotide #	NM_002755.3
UniProt Accession #	Q02750
Molecular Weight	43,439 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 1
NCBI Official Symbol	MAP2K1
NCBI Official Synonym Symbols	MEK1; MKK1; MAPKK1; PRKMK1
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 1; MEK 1; MAPKK 1; MAPK/ERK kinase 1; ERK activator kinase 1; protein kinase, mitogen-activated, kinase 1 (MAP kinase kinase 1)
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 1

UniProt Gene Name	MAP2K1
UniProt Synonym Gene Names	MEK1; PRKMK1; MAP kinase kinase 1; MAPKK 1; MKK1; MEK 1
UniProt Entry Name	MP2K1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/map2k1-recombinant-antibody-clone-9b6-immunoprecipitation-immunohistochemistry-western-blot-elisa/243994>