

CD274 recombinant antibody

CD274 Antibody

CATALOG # AAA244004	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CD274; Monoclonal Recombinant Antibody; CD274 Antibody; Programmed cell death 1 ligand 1 (PD-L1) (PDCD1 ligand 1) (Programmed death ligand 1) (B7 homolog 1) (B7-H1) (CD antigen CD274); B7H1 PDCD1L1 PDCD1LG1 PDL1; CD274 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	10D4
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for CD274 recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human PD-L1 (CD274)
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human lung cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: HepG2 whole cell lysate, K562 whole cell lysate, A549 whole cell lysate, HeLa whole cell lysate. All lanes: CD274 Antibody at 1:1000. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 34, 21, 21 kDa. Observed band size: 45 kDa.

ADDITIONAL INFORMATION

Related Product Information for CD274 recombinant antibody	Involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation and cytokine production.
Product Categories/Family for CD274 recombinant antibody	Cancer; Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	390979639
NCBI GeneID	29126
NCBI Accession #	NP_001254635.1
NCBI GenBank Nucleotide #	NM_001267706.1
UniProt Accession #	Q9NZQ7; Q14CJ2; Q2V8D5; Q66RK1; Q6WEX4; Q9NUZ5; B2RBA2; B4DU27
Molecular Weight	33,275 Da
NCBI Official Full Name	programmed cell death 1 ligand 1 isoform b
NCBI Official Synonym Full Names	CD274 molecule
NCBI Official Symbol	CD274
NCBI Official Synonym Symbols	B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1
NCBI Protein Information	programmed cell death 1 ligand 1; B7 homolog 1; CD274 antigen; PDCD1 ligand 1; programmed death ligand 1
UniProt Protein Name	Programmed cell death 1 ligand 1
UniProt Gene Name	CD274
UniProt Synonym Gene Names	B7H1; PDCD1L1; PDCD1LG1; PDL1; PD-L1; PDCD1 ligand 1; Programmed death ligand 1; B7-H1
UniProt Entry Name	PD1L1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd274-recombinant-antibody-clone-10d4-immunohistochemistry-western-blot-elisa/244004>