

SYNCRIP recombinant antibody

SYNCRIP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA244010	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SYNCRIP; Monoclonal Recombinant Antibody; SYNCRIP Antibody; Heterogeneous nuclear ribonucleoprotein Q (hnRNP Q) (Glycine- and tyrosine-rich RNA-binding protein) (GRY-RBP) (NS1-associated protein 1) (Synaptotagmin-binding; cytoplasmic RNA-interacting protein); HNRPQ NSAP1; SYNCRIP recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	9C7
Purity/Purification	Affinity-chromatography
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, PH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for SYNCRIP recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Antibody Type	Recombinant Antibody
Conjugation	Non-conjugated
Immunogen	A synthesized peptide derived from human hnRNP Q
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HeLa Cells at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeated by 0.2% TritonX-100, and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. Nuclear DNA was labeled in blue with DAPI. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Rabbit IgG ?H+L?.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for SYNCRIP recombinant antibody	Heterogenous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1, isoform 2 and isoform 3 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences. Isoform 1 is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 3 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins. Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma activation assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation; seems not to be essential for GAIT complex function.
Product Categories/Family for SYNCRIP recombinant antibody	Epigenetics and Nuclear Signaling ; Microbiology

NCBI AND UNIPROT INFORMATION

NCBI GI #	228008396
NCBI GeneID	10492
NCBI Accession #	NP_001153145.1
NCBI GenBank Nucleotide #	NM_001159673.1
UniProt Accession #	O60506 ; Q53H05 ; Q5TCG2 ; Q5TCG3 ; Q8IW78 ; Q8N599 ; Q96LC1 ; Q96LC2 ; Q9Y583 ; E1P501 ; E1P502
Molecular Weight	69,603 Da
NCBI Official Full Name	heterogeneous nuclear ribonucleoprotein Q isoform 2
NCBI Official Synonym Full Names	synaptotagmin binding, cytoplasmic RNA interacting protein
NCBI Official Symbol	SYNCRIP
NCBI Official Synonym Symbols	PP68; NSAP1; GRYRBP; HNRNPQ; HNRPQ1; GRY-RBP; hnRNP-Q
NCBI Protein Information	heterogeneous nuclear ribonucleoprotein Q; NS1-associated protein 1; glycine- and tyrosine-rich RNA-binding protein
UniProt Protein Name	Heterogeneous nuclear ribonucleoprotein Q

UniProt Gene Name	SYNCRIP
UniProt Synonym Gene Names	HNRPQ; NSAP1; hnRNP Q; GRY-RBP
UniProt Entry Name	HNRPQ_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/syncrip-recombinant-antibody-clone-9c7-immunofluorescence-immunohistochemistry-elisa/244010>