

CD40 ligand (CD40LG) Active Protein | CD40LG active protein

Recombinant Human CD40 ligand (CD40LG), partial

CATALOG #
AAA244028

HOST
Mammalian Cell


PRODUCT SYNONYMS

CD40 ligand (CD40LG); Recombinant Human CD40 ligand (CD40LG); partial; CD40-L; T-cell antigen Gp39; NF-related activation protein; TRAP; Tumor necrosis factor ligand superfamily member 5; CD154; sCD40L; CD40LG active protein


PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. Lyophilized from a 0.2 um filtered PBS, 6% Trehalose, pH7.4 Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	13-261aa, Partial
Sequence	MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYTMSNNL VTLENGKQLT VKRQGLYYIY AQVTFCNRE ASSQAPFIAS LCLKSPGRFE RILLRAANTH SSAKPCGQQS IHLGGVFELQ PGASVFVNVT DPSQVSHGTG FTSFGLLKL
Species	Human
Endotoxin	Less than 1.0EU/ug as determined by LAL method.
Biological Activity	1: Measured by its binding ability in a functional ELISA. Immobilized CD40L at 2ug/ml can bind Anti-CD40L Rabbit Monoclonal Antibody the EC50 is 2.353-3.232ng/ml.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Preparation and Storage	Store at -20 degrees C. For long-term storage, store at -20 degrees C or -80 degrees C. Store working aliquots at 4 degrees C for up to one week. Repeated freezing and thawing is not recommended.

IMAGES



Bioactivity



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for CD40LG active protein	Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:1280226, PubMed:31331973). Costimulates T-cell proliferation and cytokine production (PubMed:8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:8617933). Induces the activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching
Product Categories/Family for CD40LG active protein	Cancer

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557433
NCBI GeneID	959
NCBI Accession #	NP_000065.1
NCBI GenBank Nucleotide #	NM_000074.2
UniProt Accession #	P29965
Molecular Weight	29,274 Da
NCBI Official Full Name	CD40 ligand
NCBI Official Synonym Full Names	CD40 ligand
NCBI Official Symbol	CD40LG
NCBI Official Synonym Symbols	IGM; IMD3; TRAP; gp39; CD154; CD40L; HIGM1; T-BAM; TNFSF5; hCD40L
NCBI Protein Information	CD40 ligand; CD40-L; CD40 antigen ligand; T-cell antigen Gp39; T-B cell-activating molecule; TNF-related activation protein; tumor necrosis factor (ligand) superfamily member 5
UniProt Protein Name	CD40 ligand
UniProt Gene Name	CD40LG
UniProt Synonym Gene Names	CD40L; TNFSF5; TRAP; CD40-L; TRAP
UniProt Entry Name	CD40L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mammalian-cell-active-protein/cd40-ligand-cd40lg/244028>