

Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B) Active Protein | TNFRSF1B active protein

Recombinant Human Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B), partial

CATALOG #
AAA244030


HOST
Mammalian Cell

PRODUCT SYNONYMS


Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B); Recombinant Human Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B); partial; Tumor necrosis factor receptor 2; TNF-R2; Tumor necrosis factor receptor type II; TNF-RII; TNFR-II; p75; p80 TNF-alpha receptor; CD120b; Etanercept; TBP-2; TBPII; TNFRSF1B active protein

PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. Lyophilized from a 0.2 um filtered PBS, 6% Trehalose, pH7.4 Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	23-257aa, Partial
Sequence	LPAQVAFPTY APEPGSTCRL REYYDQTAQM CCSKCSPGQH AKVFCTKTS TVCDSCEDST YTQLWNWVPE CLSCGSRCSS DQVETQACTR EQNRICTCRP GWYCALSKQE GCRLCAPLRK CRPGFGVARP GTETSDVVCK PCAPGTFSNT TSSTDICRPH QICNVVAIPG NASMDAVCTS TSPTRSMAPG AVHLPQPVST RSQHTQPTPE PSTAPSTSFL LPMGPPSPAE GSTGD
Species	Human
Endotoxin	Less than 1.0EU/ug as determined by LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized TNF-alpha at 5ug/ml can bind human TNFR2, the EC50 of human TNFR2 protein is 1.162-1.481ng/ml.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Preparation and Storage	Store at -20 degrees C. For long-term storage, store at -20 degrees C or -80 degrees C. Store working aliquots at 4 degrees C for up to one week. Repeated freezing and thawing is not recommended.



Bioactivity



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for TNFRSF1B active protein	Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.
Product Categories/Family for TNFRSF1B active protein	Cancer

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507577
NCBI GeneID	7133
NCBI Accession #	NP_001057.1
NCBI GenBank Nucleotide #	NM_001066.2
UniProt Accession #	P20333; Q16042; Q6YI29; Q9UIH1; B1AJZ3
Molecular Weight	48,291 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 1B
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 1B
NCBI Official Symbol	TNFRSF1B
NCBI Official Synonym Symbols	p75; TBPII; TNFBR; TNFR2; CD120b; TNFR1B; TNFR80; TNF-R75; p75TNFR; TNF-R-II
NCBI Protein Information	tumor necrosis factor receptor superfamily member 1B; TNF-R2; TNF-R-II; p75 TNF receptor; p80 TNF-alpha receptor; soluble TNFR1B variant 1; tumor necrosis factor receptor 2; tumor necrosis factor beta receptor; tumor necrosis factor receptor type II; tumor
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 1B
UniProt Gene Name	TNFRSF1B
UniProt Synonym Gene Names	TNFBR; TNFR2; TNF-R2; TNF-R-II; TNFR-II
UniProt Entry Name	TNR1B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/mammalian-cell-active-protein/tumor-necrosis-factor-receptor-superfamily-member-1b-tnfrsf1b/244030>