

Rabbit anti-Human CLTC Polyclonal Antibody | anti-CLTC antibody

Rabbit anti-Human CLTC Polyclonal Antibody

CATALOG # AAA244060	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CLTC; Polyclonal Antibody; Rabbit anti-Human CLTC Polyclonal Antibody; Clathrin heavy chain 1 (Clathrin heavy chain on chromosome 17) (CLH-17); CLH17 CLTCL2 KIAA0034; CLTC Antibody; anti-CLTC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-CLTC antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Clathrin heavy chain 1 protein (851-990aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of diluted at 1:200 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for anti-CLTC antibody	Clathrin is the major protein of the polyhedral coat of coated pits and vesicles. Two different adapter protein complexes link the clathrin lattice either to the plasma membrane or to the trans-Golgi network. Acts as component of the TACC3/ch-TOG/clathrin complex proposed to contribute to stabilization of kinetochore fibers of the mitotic spindle by acting as inter-microtubule bridge (PubMed:15858577, PubMed:16968737, PubMed:21297582). The TACC3/ch-TOG/clathrin complex is required for the maintenance of kinetochore fiber tension (PubMed:23532825). Plays a role in early autophagosome formation (PubMed:20639872).
Product Categories/Family for anti-CLTC antibody	Others

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758012
NCBI GeneID	1213
NCBI Accession #	NP_004850.1
NCBI GenBank Nucleotide #	NM_004859.3
UniProt Accession #	Q00610; Q6N0A0; Q86TF2; D3DU00
Molecular Weight	191615 MW
NCBI Official Full Name	clathrin heavy chain 1 isoform 1
NCBI Official Synonym Full Names	clathrin heavy chain
NCBI Official Symbol	CLTC
NCBI Official Synonym Symbols	Hc; CHC; CHC17; CLH-17; CLTCL2
NCBI Protein Information	clathrin heavy chain 1
UniProt Protein Name	Clathrin heavy chain 1
UniProt Gene Name	CLTC
UniProt Synonym Gene Names	CLH17; CLTCL2; KIAA0034; CLH-17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cltc-antibody-immunohistochemistry-elisa/244060>