

Rabbit anti-Human EIF5A Polyclonal Antibody | anti-EIF5A antibody

Rabbit anti-Human EIF5A Polyclonal Antibody

CATALOG # AAA244062	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

EIF5A; Polyclonal Antibody; Rabbit anti-Human EIF5A Polyclonal Antibody; Eukaryotic translation initiation factor 5A-1 (eIF-5A-1) (eIF-5A1) (Eukaryotic initiation factor 5A isoform 1) (eIF-5A) (Rev-binding factor) (eIF-4D); EIF5A Antibody; anti-EIF5A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-EIF5A antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Eukaryotic translation initiation factor 5A-1 protein (2-154aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of diluted at 1:200 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of diluted at 1:200 and staining in paraffin-embedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for anti-EIF5A antibody	mRNA-binding protein involved in translation elongation. Has an important function at the level of mRNA turnover, probably acting downstream of decapping. Involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity. With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis. Regulates also TNF-alpha-mediated apoptosis. Mediates effects of polyamines on neuronal process extension and survival. May play an important role in brain development and function, and in skeletal muscle stem cell differentiation. Also described as a cellular cofactor of human T-cell leukemia virus type I (HTLV-1) Rex protein and of human immunodeficiency virus type 1 (HIV-1) Rev protein, essential for mRNA export of retroviral transcripts.
Product Categories/Family for anti-EIF5A antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	54037409
NCBI GeneID	1984
UniProt Accession #	P63241; P10159; Q16182; Q7L7L3; Q7Z4L1; Q9D0G2; A8K9A0; D3DTP2; AAB29229
Molecular Weight	154
NCBI Official Full Name	Eukaryotic translation initiation factor 5A-1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 5A
NCBI Official Symbol	EIF5A
NCBI Official Synonym Symbols	EIF-5A; EIF5A1; eIF5AI
NCBI Protein Information	eukaryotic translation initiation factor 5A-1; eIF-4D; eIF-5A1; eIF-5A-1; rev-binding factor; eukaryotic initiation factor 5A
UniProt Protein Name	Eukaryotic translation initiation factor 5A-1
UniProt Gene Name	EIF5A
UniProt Synonym Gene Names	eIF-5A-1; eIF-5A1; eIF-5A
UniProt Entry Name	IF5A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/eif5a-antibody-immunohistochemistry-elisa/244062>