

Rabbit anti-Human HIST1H2AG Polyclonal Antibody | anti-HIST1H2AG antibody

Rabbit anti-Human HIST1H2AG Polyclonal Antibody

CATALOG # AAA244072	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

HIST1H2AG; Polyclonal Antibody; Rabbit anti-Human HIST1H2AG Polyclonal Antibody; Histone H2A type 1 (H2A.1) (Histone H2A/ptI); HIST1H2AI; HIST1H2AK; HIST1H2AL; HIST1H2AM; H2AFP; H2AFC; H2AFD; H2AFI; H2AFN; HIST1H2AG (Ab-9) Antibody; anti-HIST1H2AG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-HIST1H2AG antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Synthesized peptide derived from Human Histone H2A type 1 protein (2-14aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of diluted at 1:10 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of diluted at 1:10 and staining in paraffin-embedded human glioma cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H2AG antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H2AG antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504239
NCBI GeneID	8329
NCBI Accession #	NP_003500.1
NCBI GenBank Nucleotide #	NM_003509.2
UniProt Accession #	P0C0S8; P02261; Q2M1R2; Q76PA6
Molecular Weight	14,091 Da
NCBI Official Full Name	histone H2A type 1
NCBI Official Synonym Full Names	histone cluster 1 H2A family member i
NCBI Official Symbol	HIST1H2AI
NCBI Official Synonym Symbols	H2A/c; H2AFC
NCBI Protein Information	histone H2A type 1
UniProt Protein Name	Histone H2A type 1
UniProt Gene Name	HIST1H2AG
UniProt Synonym Gene Names	H2AFP; H2A.1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hist1h2ag-antibody-immunohistochemistry-elisa/244072>