

Rabbit anti-Human ICAM3 Polyclonal Antibody | anti-ICAM3 antibody

Rabbit anti-Human ICAM3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA244093	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

ICAM3; Polyclonal Antibody; Rabbit anti-Human ICAM3 Polyclonal Antibody; Intercellular adhesion molecule 3 (ICAM-3) (CDw50) (ICAM-R) (CD antigen CD50); ICAM3 Antibody; anti-ICAM3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-ICAM3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Intercellular adhesion molecule 3 protein (200-547aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of PC3 cells with at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of diluted at 1:100 and staining in paraffin-embedded Human spleen tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



WB (Western Blot)

Western Blot Positive WB detected in: L02 whole cell lysate, PC-3 whole cell lysate, U87 whole cell lysate, HepG2 whole cell lysate
 All lanes: ICAM3 antibody at 1:2000
 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution
 Predicted band size: 60 kDa
 Observed band size: 60 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-ICAM3 antibody	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2) (PubMed:1448173). ICAM3 is also a ligand for integrin alpha-D/beta-2. In association with integrin alpha-L/beta-2, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:23775590).
Product Categories/Family for anti-ICAM3 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	167466207
NCBI GeneID	3385
NCBI Accession #	NP_002153.2
NCBI GenBank Nucleotide #	NM_002162.3
UniProt Accession #	P32942 ; Q6PD68
Molecular Weight	59,541 Da
NCBI Official Full Name	intercellular adhesion molecule 3
NCBI Official Synonym Full Names	intercellular adhesion molecule 3
NCBI Official Symbol	ICAM3
NCBI Official Synonym Symbols	CD50; CDW50; ICAM-R
NCBI Protein Information	intercellular adhesion molecule 3; ICAM-3; intercellular adhesion molecule-3
UniProt Protein Name	Intercellular adhesion molecule 3
UniProt Gene Name	ICAM3
UniProt Synonym Gene Names	ICAM-3
UniProt Entry Name	ICAM3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.