

Rabbit anti-Human INTS10 Polyclonal Antibody | anti-INTS10 antibody

Rabbit anti-Human INTS10 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA244094	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

INTS10; Polyclonal Antibody; Rabbit anti-Human INTS10 Polyclonal Antibody; Integrator complex subunit 10 (Int10); C8orf35; INTS10 Antibody; anti-INTS10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-INTS10 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Integrator complex subunit 10 protein (51-342aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of U251 cells with at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of diluted at 1:300 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for anti-INTS10 antibody	Component of the Integrator (INT) complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes (Probable). May be not involved in the recruitment of cytoplasmic dynein to the nuclear envelope by different components of the INT complex (PubMed:23904267).
Product Categories/Family for anti-INTS10 antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	83582809
NCBI GeneID	55174
NCBI Accession #	NP_060612.2
NCBI GenBank Nucleotide #	NM_018142.2
UniProt Accession #	Q9NVR2; Q6IA93; Q7L538; Q7L8C8; Q9H3W8
Molecular Weight	82,236 Da
NCBI Official Full Name	integrator complex subunit 10
NCBI Official Synonym Full Names	integrator complex subunit 10
NCBI Official Symbol	INTS10
NCBI Official Synonym Symbols	INT10; C8orf35
NCBI Protein Information	integrator complex subunit 10
UniProt Protein Name	Integrator complex subunit 10
UniProt Gene Name	INTS10
UniProt Synonym Gene Names	C8orf35; Int10
UniProt Entry Name	INT10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ints10-antibody-immunofluorescence-immunohistochemistry-elisa/244094>