

Mouse ATF-3 (Activating Transcription Factor 3) ELISA Kit | ATF-3 elisa kit

Mouse ATF-3 (Activating Transcription Factor 3) ELISA Kit

CATALOG #
AAA248844

REACTIVITY
Mouse


PRODUCT SYNONYMS

ATF-3 (Activating Transcription Factor 3); Mouse ATF-3 (Activating Transcription Factor 3) ELISA Kit; ATF-3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of ATF-3. No significant cross-reactivity or interference between ATF-3 and analogues was observed.
Sequence Length	181
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	15.625-1000pg/ml
Sensitivity	9.375pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ATF-3 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ATF-3 were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ATF-3 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-ATF-3 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti- ATF-3 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the ATF-3 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of ATF-3 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	442422
NCBI GeneID	467
Molecular Weight	14,314 Da
NCBI Official Full Name	activating transcription factor 3
NCBI Official Synonym Full Names	activating transcription factor 3
NCBI Official Symbol	ATF3
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-3
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-3
UniProt Gene Name	ATF3
UniProt Synonym Gene Names	cAMP-dependent transcription factor ATF-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/atf-3-activating-transcription-factor-3/248844>