

Bovine CD14 (Monocyte differentiation antigen CD14) ELISA Kit | CD14 elisa kit

Bovine CD14 (Monocyte differentiation antigen CD14) ELISA Kit

CATALOG #
AAA248906

REACTIVITY
Bovine

PRODUCT SYNONYMS

CD14 (Monocyte differentiation antigen CD14); Bovine CD14 (Monocyte differentiation antigen CD14) ELISA Kit; CD14 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Bovine
Specificity	This assay has high sensitivity and excellent specificity for detection of CD109. No significant cross-reactivity or interference between CD109 and analogues was observed.
Sequence Length	375
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	15.625-1000ng/ml
Sensitivity	9.375ng/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CD109 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CD109 were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve

Results of a typical standard operation of a CD14 ELISA Kit are listed below. This standard curve was generated at our lab for demonstration purpose only. Users shall obtain standard curve as per experiment by themselves. (N/A=not applicable)

X	ng/ml	0	15.625	31.25	62.5	125	250	500	1000
Y	OD450	0.075	0.098	0.185	0.267	0.443	0.77	1.375	2.06

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD14 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-CD109 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-CD109 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the CD109 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of CD109 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557417
NCBI GeneID	929
NCBI Accession #	NP_000582.1
NCBI GenBank Nucleotide #	NM_000591.4
UniProt Accession #	P08571; Q53XT5; Q96FR6; Q96L99; Q9UNS3
Molecular Weight	40,076 Da
NCBI Official Full Name	monocyte differentiation antigen CD14
NCBI Official Synonym Full Names	CD14 molecule
NCBI Official Symbol	CD14
NCBI Protein Information	monocyte differentiation antigen CD14
UniProt Protein Name	Monocyte differentiation antigen CD14
UniProt Gene Name	CD14

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/bovine-ovine-elisa-kits/cd14-monocyte-differentiation-antigen-cd14/248906>