

Human STOM (Stomatin) ELISA Kit | STOM elisa kit

Human STOM (Stomatin) ELISA Kit

CATALOG #
AAA248934

REACTIVITY
Human

PRODUCT SYNONYMS

STOM (Stomatin); Human STOM (Stomatin) ELISA Kit; STOM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of STMN1. No significant cross-reactivity or interference between STMN1 and analogues was observed.
Sequence Length	288
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	0.156-10ng/ml
Sensitivity	0.094ng/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level STMN1 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level STMN1 were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve

Results of a typical standard operation of a STOM ELISA Kit are listed below. This standard curve was generated at our lab for demonstration purpose only. Users shall obtain standard curve as per experiment by themselves. (N/A-not applicable)

X	ng/ml	0	0.156	0.312	0.625	1.25	2.5	5	10
Y	OD450	0.077	0.129	0.176	0.268	0.442	0.775	1.417	1.983

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for STOM elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-STMN1 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-STMN1 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the STMN1 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of STMN1 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	14715077
NCBI GeneID	2040
Molecular Weight	13,475 Da
NCBI Official Full Name	Stomatin
NCBI Official Synonym Full Names	stomatin
NCBI Official Symbol	STOM
NCBI Official Synonym Symbols	BND7; EPB7; EPB72
NCBI Protein Information	erythrocyte band 7 integral membrane protein
UniProt Protein Name	Erythrocyte band 7 integral membrane protein
UniProt Gene Name	STOM
UniProt Synonym Gene Names	BND7; EPB72

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/stom-stomatin/248934>