

Rabbit TGF- β 3 (Transforming Growth Factor beta3) ELISA Kit | TGF- β 3 elisa kit

Rabbit TGF- β 3 (Transforming Growth Factor beta3) ELISA Kit

CATALOG #
AAA249020

REACTIVITY
Rabbit

PRODUCT SYNONYMS

TGF- β 3 (Transforming Growth Factor beta3); Rabbit TGF- β 3 (Transforming Growth Factor beta3) ELISA Kit; TGF- β 3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rabbit
Specificity	This assay has high sensitivity and excellent specificity for detection of TGF β 1. No significant cross-reactivity or interference between TGF β 1 and analogues was observed.
Sequence Length	412
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	15.625-1000pg/ml
Sensitivity	9.375pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TGF β 1 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TGF β 1 were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve

Results of a typical standard operation of a TGF- β 3 ELISA Kit are listed below. This standard curve was generated at our lab for demonstration purpose only. Users shall obtain standard curve as per experiment by themselves. (N/A=not applicable)

X	pg/ml	0	15.625	31.25	62.5	125	250	500	1000
Y	OD450	0.049	0.102	0.171	0.283	0.515	0.935	1.611	2.231

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TGF- β 3 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-TGF β 1 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-TGF β 1 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the TGF β 1 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of TGF β 1 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	6981650
NCBI GeneID	25717
NCBI Accession #	NP_037306.1
NCBI GenBank Nucleotide #	NM_013174.2
UniProt Accession #	Q07258; Q56A31
Molecular Weight	47,116 Da
NCBI Official Full Name	transforming growth factor beta-3 proprotein preproprotein
NCBI Official Synonym Full Names	transforming growth factor, beta 3
NCBI Official Symbol	Tgfb3
NCBI Official Synonym Symbols	TGF-B3
NCBI Protein Information	transforming growth factor beta-3 proprotein; transforming growth factor beta-3
UniProt Protein Name	Transforming growth factor beta-3 proprotein
UniProt Gene Name	Tgfb3
UniProt Synonym Gene Names	Tgf-b3; LAP; TGF-beta-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-elisa-kits/tgf-b3-transforming-growth-factor-beta3/249020>