

Mouse anti-Human ATP2C1 Monoclonal Antibody | anti-ATP2C1 antibody

ATP2C1 (KIAA1347, PMR1L, Calcium-transporting ATPase Type 2C Member 1, ATP-dependent Ca(2+) Pump PMR1) (FITC)

CATALOG # AAA25031	HOST Mouse	REACTIVITY Human	APPLICATIONS ELISA, Immunofluorescence, Immunoprecipitation, Western Blot
------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

ATP2C1; Monoclonal Antibody; ATP2C1 (KIAA1347; PMR1L; Calcium-transporting ATPase Type 2C Member 1; ATP-dependent Ca(2+) Pump PMR1) (FITC); anti-ATP2C1 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1, lambda
Clone Number	2G1
Specificity	Recognizes human ATP2C1.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with Fluorescein isothiocyanate (FITC).
Applicable Applications for anti-ATP2C1 antibody	ELISA, IF (Immunofluorescence), IP (Immunoprecipitation), WB (Western Blot)
Application Notes	IF: 10ug/ml Applications are based on unconjugated antibody.
Immunogen	Partial recombinant corresponding to aa119-269 from human ATP2C1 (AAH28139) with GST tag. MW of the GST tag alone is 26kD.
Immunogen Sequence	VQEYRSEKSLEELSKLVPPECHCVREGKLEHTLARDLVPGDTVCLSVGDRVPADRLRFEAVDLSIDESSLTGETTPCSKVTPAQPAATNGDLASRSNIAFM
Conjugate	FITC
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C. Thaw and use immediately. Do not freeze and thaw repeatedly. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Caution: FITC conjugates are sensitive to light and may lose activity upon repeated freezing and thawing and prior to removing the cap.

IMAGES



WB (Western Blot)

ATP2C1 monoclonal antibody Western Blot analysis of ATP2C1 expression in HeLa.




WB (Western Blot)

Western blot analysis of ATP2C1 over-expressed 293 cell line, cotransfected with ATP2C1 Validated Chimera RNAi (Lane 2) or non-transfected control (Lane 1). Blot probed with ATP2C1 monoclonal antibody. GAPDH (36.1kD) used as specificity and loading control.




Application Data

Detection limit for recombinant GST tagged ATP2C1 is ~0.03ng/ml as a capture antibody.




IP (Immunoprecipitation)

Immunoprecipitation of ATP2C1 transfected lysate using ATP2C1 monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with ATP2C1 rabbit polyclonal antibody.



IF (Immunofluorescence)

Immunofluorescence of monoclonal antibody to ATP2C1 on HeLa cell. [antibody concentration 10ug/ml]



WB (Western Blot)

Western Blot analysis of ATP2C1 expression in transfected 293T cell line by ATP2C1 monoclonal antibody. Lane 1: ATP2C1 transfected lysate (100.6kD). Lane 2: Non-transfected lysate.

ADDITIONAL INFORMATION

Product Categories/Family for anti-ATP2C1 antibody

Antibodies; Ion Channel

NCBI AND UNIPROT INFORMATION

NCBI GI #	20380102
NCBI GeneID	27032
Molecular Weight	104,032 Da
NCBI Official Full Name	Homo sapiens ATPase, Ca ⁺⁺ transporting, type 2C, member 1, mRNA
NCBI Official Synonym Full Names	ATPase secretory pathway Ca ²⁺ transporting 1
NCBI Official Symbol	ATP2C1
NCBI Official Synonym Symbols	HHD; BCPM; PMR1; SPCA1; hSPCA1; ATP2C1A
NCBI Protein Information	calcium-transporting ATPase type 2C member 1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/atp2c1-antibody-clone-2g1-immunofluorescence-immunoprecipitation-western-blot-elisa/25031>