

Mouse Ubiquitin Monoclonal Antibody | anti-UBB antibody

Ubiquitin Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA253941	Mouse	Human, Mouse, Rat, Bovine	ELISA, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS


Ubiquitin; Monoclonal Antibody; Ubiquitin Antibody; Mouse Anti-Bovine Ubiquitin Monoclonal IgG2a Kappa; Clone 6C11-B3; Polyubiquitin B Antibody; RPS27A Antibody; UBA52 Antibody; UBB Antibody; UBC Antibody; ubiquitin B Antibody; anti-UBB antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat, Bovine
Clonality	Monoclonal
Isotype	IgG2a Kappa
Clone Number	6C11-B3
Specificity	Detects ~10kDa.
Purity/Purification	Protein G Purified
Form/Format	PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide
Concentration	1 mg/ml (varies by lot)
Sequence Length	305
Applicable Applications for anti-UBB antibody	ELISA, IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	Native bovine ubiquitin, conjugated to KLH
Immunogen Species	Bovine
Conjugate	Unconjugated
Cellular Localization	Cytoplasm, Nucleus
Certificate of Analysis	1 ug/ml of SMC-171 was sufficient for detection of ubiquitin in 10 ug of Hela Lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Preparation and Storage	Store at -20 degree C.



ICC (Immunocytochemistry)
Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Ubiquitin Monoclonal Antibody, Clone 6C11-B3 . Tissue: Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-Ubiquitin Monoclonal Antibody at 1:100 for 12 hours at 4 degree C. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Diffuse nuclear and cytoplasmic staining. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Ubiquitin Antibody. (C) Composite.



WB (Western Blot)
Western Blot analysis of Human Cell lysates showing detection of Ubiquitin protein using Mouse Anti-Ubiquitin Monoclonal Antibody, Clone 6C11-B3 . Load: 15 ug. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Ubiquitin Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



ICC (Immunocytochemistry)
Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Ubiquitin Monoclonal Antibody, Clone 6C11-B3 . Tissue: Cervical cancer cell line (HeLa). Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-Ubiquitin Monoclonal Antibody at 1:100 for 12 hours at 4 degree C. Secondary Antibody: APC Goat Anti-Mouse (red) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Diffuse nuclear and cytoplasmic staining. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Ubiquitin Antibody. (C) Composite.

ADDITIONAL INFORMATION

Related Product Information for anti-UBB antibody	Ubiquitin is a small protein that occurs in all eukaryotic cells. The ubiquitin protein itself consists of 76 amino acids and has a molecular mass of about 8.5kDa. Key features include its C-terminal tail and the 7 Lys residues. It is highly conserved among eukaryotic species: Human and yeast ubiquitin share 96% sequence identity (1). The main function of Ubiquitin is to clear abnormal, foreign and improperly folded proteins by targeting them for degradation by the 26S proteasome (2). Ubiquitination represents an essential cellular process affected by a multi-enzyme cascade involving classes of enzymes known as ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s or Ubcs) and ubiquitin-protein ligases (E3s). Ubiquitin is activated in a two-step reaction by an E1 ubiquitin-activating enzyme in a process requiring ATP as an energy source. The initial step involves production of an ubiquitin-adenylate intermediate. The second step transfers ubiquitin to the E1 active site cysteine residue, with release of AMP. This step results in a thioester linkage between the C-terminal carboxyl group of ubiquitin and the E1 cysteine sulfhydryl group. The third step is a transfer of ubiquitin from E1 to the active site cysteine of a ubiquitin-conjugating enzyme E2 via a trans(thio)esterification reaction. And the final step of the ubiquitylation cascade creates an isopeptide bond between a lysine of the target protein and the C-terminal glycine of ubiquitin. In general, this step requires the activity of one of the hundreds of E3 ubiquitin-protein ligases (often termed simply ubiquitin ligase). E3 enzymes function as the substrate recognition modules of the system and are capable of interaction with both E2 and substrate(2, 3). Ubiquitination also participates in the internalization and degradation of plasma membrane proteins such as some of the TCR subunits while still ER-membrane associated (4). Ubiquitin also plays a role in regulating signal transduction cascades through the elimination inhibitory proteins, such as IkappaBalpha and p27 (5).
Product Categories/Family for anti-UBB antibody	Cell Signaling ; Post-translational Modifications ; Ubiquitination ; Neuroscience ; Neurodegeneration ; Alzheimer's Disease
References	1. Wilkinson K.D. (1995) Annu. Rev. Nutr. 15:161-189. 2. Bonifacino J.S., et al. (1998) Annu Rev Cell Dev Biol. 14: 19-57. 3. Boston Biochem: "Ubiquitin Proteasome Pathway Overview". 4. Yang M., et al. (1998) J Exp Med. 187: 1835-1846. 5. Chen Z.J., et al. (1996) Cell 84: 853-862.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27806505
NCBI GeneID	281370
NCBI Accession #	NP_776558
NCBI GenBank Nucleotide #	NM_174133.2
UniProt Accession #	P0CG53
NCBI Official Full Name	polyubiquitin-B
NCBI Official Symbol	UBB
NCBI Protein Information	polyubiquitin-B
UniProt Protein Name	Polyubiquitin-B
UniProt Gene Name	UBB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/ubiquitin-antibody-clone-6c11-b3-immunofluorescence-immunocytochemistry-western-blot-elisa/253941>