

Rabbit anti-Human EphB2 Polyclonal Antibody | anti-EphB2 antibody

Anti-EphB2 Antibody, Rabbit Polyclonal

CATALOG # AAA254133	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

EphB2; Polyclonal Antibody; Anti-EphB2 Antibody; Rabbit Polyclonal; EPHB2 antibody; Rabbit PAb; Antigen Affinity Purified; EPH receptor B2; Anti-CAPB Antibody; Anti-DRT Antibody; Anti-EK5 Antibody; Anti-EPHT3 Antibody; Anti-ERK Antibody; Anti-Hek5 Antibody; Anti-PCBC Antibody; Anti-Tyro5 Antibody; anti-EphB2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human EPHB2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-EphB2 antibody	IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to the N-terminus of the human EPHB2
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with A synthetic peptide corresponding to the N-terminus of the human EPHB2., and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunochemical staining of human EPHB2 in human pancreas with rabbit polyclonal antibody (1:300, formalin-fixed paraffin embedded sections).


IHC (Immunohistochemistry)

Immunochemical staining of human EPHB2 in human placenta with rabbit polyclonal antibody (1:300, formalin-fixed paraffin embedded sections).



IHC (Immunohistochemistry)
 Immunochemical staining of human EPHB2 in human skeletal muscle with rabbit polyclonal antibody (1:300, formalin-fixed paraffin embedded sections).

ADDITIONAL INFORMATION

Related Product Information for anti-EphB2 antibody	Ephrin type-B receptor 2, also known as EphB2, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. EphB2 receptor tyrosine kinase phosphorylates syndecan-2 and that this phosphorylation event is crucial for syndecan-2 clustering and spine formation. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity ⁶ . Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). Ligand-mediated activation of Ephs induces various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer. EphB receptor tyrosine kinases are enriched at synapses, suggesting that these receptors play a role in synapse formation or function. We find that EphrinB binding to EphB induces a direct interaction of EphB with NMDA-type glutamate receptors. This interaction occurs at the cell surface and is mediated by the extracellular regions of the two receptors, but does not require the kinase activity of EphB.
Product Categories/Family for anti-EphB2 antibody	Actin Dynamics Signaling Pathway
References	Zisch AH, et al. (1998) Complex formation between EphB2 and Src requires phosphorylation of tyrosine 611 in the EphB2 juxtamembrane region. <i>Oncogene</i> . 16 (20): 2657-70. Yu HH, et al. (2001) Multiple signaling interactions of Abl and Arg kinases with the EphB2 receptor. <i>Oncogene</i> . 20 (30): 3995-4006. Zisch AH, et al. (2000) Replacing two conserved tyrosines of the EphB2 receptor with glutamic acid prevents binding of SH2 domains without abrogating kinase activity and biological responses. <i>Oncogene</i> . 19 (2): 177-87.

NCBI AND UNIPROT INFORMATION

NCBI GI #	21396504
NCBI GeneID	2048
NCBI Accession #	NP_004433.2
NCBI GenBank Nucleotide #	NM_004442.6
UniProt Accession #	P29323 ; O43477 ; Q5T0U6 ; Q5T0U7 ; Q5T0U8
Molecular Weight	117,493 Da
NCBI Official Full Name	ephrin type-B receptor 2 isoform 2
NCBI Official Synonym Full Names	EPH receptor B2
NCBI Official Symbol	EPHB2
NCBI Official Synonym Symbols	DRT; EK5; ERK; CAPB; Hek5; PCBC; EPHT3; Tyro5

NCBI Protein Information	ephrin type-B receptor 2; EPH-like kinase 5; eph tyrosine kinase 3; elk-related tyrosine kinase; protein-tyrosine kinase HEK5; tyrosine-protein kinase TYRO5; renal carcinoma antigen NY-REN-47; tyrosine-protein kinase receptor EPH-3; developmentally-regula
UniProt Protein Name	Ephrin type-B receptor 2
UniProt Gene Name	EPHB2
UniProt Synonym Gene Names	DRT; EPHT3; EPTH3; ERK; HEK5; TYRO5; EK5; hEK5
UniProt Entry Name	EPHB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ephb2-antibody-immunohistochemistry-elisa/254133>