

Rabbit anti-Human CD45 Monoclonal Antibody | anti-PTPRC antibody

Recombinant Anti-CD45 Antibody, Rabbit Monoclonal

CATALOG # AAA254199	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD45; Monoclonal Antibody; Recombinant Anti-CD45 Antibody; Rabbit Monoclonal; CD45/PTPRC Antibody; Rabbit MAb; protein tyrosine phosphatase; receptor type; C; Anti-B220 Antibody; Anti-CD45 Antibody; Anti-CD45R Antibody; Anti-GP180 Antibody; Anti-L-CA Antibody; Anti-LCA Antibody; Anti-LY5 Antibody; Anti-T200 Antibody; anti-PTPRC antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	145
Specificity	Human CD45/PTPRC
Purity/Purification	Protein A
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-PTPRC antibody	IHC (Immunohistochemistry)
Antibody Type	Recombinant Antibody
Immunogen	A synthetic peptide corresponding to the C-terminus of the Human CD45/PTPRC
Conjugation	Unconjugated
Preparation	This antibody was obtained from a rabbit immunized with a synthetic peptide corresponding to the C-terminus of the Human CD45/PTPRC.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Immunochemical staining of human LCA in human lymphoma with rabbit monoclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)
Immunochemical staining of human LCA in human spleen with rabbit monoclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-PTPRC antibody	The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexes or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.
Product Categories/Family for anti-PTPRC antibody	Phosphatases and Regulators; B Cell Receptor Signaling Pathway
References	Zola H, et al. (2007) CD molecules 2006-human cell differentiation molecules. J Immunol Methods. 318 (1-2): 1-5.Ho IC, et al. (2009) GATA3 and the T-cell lineage: essential functions before and after T-helper-2-cell differentiation. Nat Rev Immunol. 9 (2): 125-35.Matesanz-Isabel J, et al. (2011) New B-cell CD molecules. Immunology Letters.134 (2): 104-12.Irie-Sasaki J, et al. (2001) CD45 is a JAK phosphatase and negatively regulates cytokine receptor signaling. Nature. 409: 349-54.

NCBI AND UNIPROT INFORMATION

NCBI GI #	392307007
NCBI GeneID	5788
NCBI Accession #	NP_002829.3
NCBI GenBank Nucleotide #	NM_002838.4
UniProt Accession #	P08575; Q16614; Q9H0Y6; A8K7W6
Molecular Weight	147,254 Da
NCBI Official Full Name	receptor-type tyrosine-protein phosphatase C isoform 1
NCBI Official Synonym Full Names	protein tyrosine phosphatase, receptor type, C
NCBI Official Symbol	PTPRC
NCBI Official Synonym Symbols	LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180

NCBI Protein Information	receptor-type tyrosine-protein phosphatase C; CD45 antigen; T200 glycoprotein; T200 leukocyte common antigen; protein tyrosine phosphatase, receptor type, c polypeptide
UniProt Protein Name	Receptor-type tyrosine-protein phosphatase C
UniProt Gene Name	PTPRC
UniProt Synonym Gene Names	CD45; L-CA
UniProt Entry Name	PTPRC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd45-antibody-clone-145-immunohistochemistry-elisa/254199>