

Rabbit anti-Human CAMKIV Polyclonal Antibody | anti-CAMK4 antibody

Anti-CAMKIV Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA254322	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

CAMKIV; Polyclonal Antibody; Anti-CAMKIV Antibody; Rabbit Polyclonal; CAMKIV/CAMK4 Antibody; Rabbit PAb; Antigen Affinity Purified; calcium/calmodulin-dependent protein kinase IV; Anti-caMK Antibody; Anti-CaMK-GR Antibody; Anti-CaMKIV Antibody; Anti-IV Antibody; anti-CAMK4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CAMKIV/CAMK4
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CAMK4 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the C-terminus of the Human CAMKIV/CAMK4
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human CAMKIV/CAMK4, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



WB (Western Blot)

Anti-CAMK4 rabbit polyclonal antibody at 1:500 dilution. Lane A: HeLa Whole Cell Lysate. Lane B: HEK293 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution. Developed using the Odyssey technique. Performed under reducing conditions. Predicted band size: 52 kDa. Observed band size: 52 kDa.



IF (Immunofluorescence)

Immunofluorescence staining of CAMK4 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human CAMK4 polyclonal antibody (1:1000) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 594-conjugated Goat Anti-rabbit IgG secondary antibody (red). Positive staining was localized to cytoplasm and nucleus.

ADDITIONAL INFORMATION

Related Product Information for anti-CAMK4 antibody	Ca2+/ calmodulin-dependent protein kinase 4 (CAMKIV) belongs to the serine/threonine protein kinase family, and to the Ca2+/calmodulin-dependent protein kinase subfamily which is widely recognized as an essential enzyme implicated in the phosphoinositide amplification cascade. Ca2+/calmodulin dependent protein kinase (CAMK) can be activated by the intracellular increased Ca2+ and then apt to combine with the target protein. Ca2+/ calmodulin-dependent protein kinase 4 (CAMKIV) is a multifunctional CaM-dependent kinase protein with limited tissue distribution, that has been implicated in transcriptional regulation in lymphocytes, neurons and male germ cells. All of the isoforms of this family, including myosin light chain kinase, phosphorylase kinase, CaMK1, CaMKII α and CaMKIV have EF-hand structure.
Product Categories/Family for anti-CAMK4 antibody	EGFR Signaling Pathway
References	Feliciano DM, et al. (2009) Repression of Ca2+/calmodulin-dependent protein kinase IV signaling accelerates retinoic acid-induced differentiation of human neuroblastoma cells. J Biol Chem. 284 (39): 26466-81. Zhao X, et al. (2001). The modular nature of histone deacetylase HDAC4 confers phosphorylation-dependent intracellular trafficking. J Biol Chem. 276 (37): 35042-8. Racioppi L, et al. (2008) Calcium/calmodulin-dependent kinase IV in immune and inflammatory responses: novel routes for an ancient traveller.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502557
NCBI GeneID	814
NCBI Accession #	NP_001735.1
NCBI GenBank Nucleotide #	NM_001744.4
UniProt Accession #	Q16566 ; D3DSZ7
Molecular Weight	51,926 Da
NCBI Official Full Name	calcium/calmodulin-dependent protein kinase type IV
NCBI Official Synonym Full Names	calcium/calmodulin-dependent protein kinase IV
NCBI Official Symbol	CAMK4
NCBI Official Synonym Symbols	IV; caMK; CaMK IV; CaMK-GR
NCBI Protein Information	calcium/calmodulin-dependent protein kinase type IV; CAM kinase IV; caM kinase-GR; CAM kinase- GR; brain Ca++-calmodulin-dependent protein kinase type IV; brain Ca(2+)-calmodulin-dependent protein kinase type IV; calcium/calmodulin-dependent protein kinase
UniProt Protein Name	Calcium/calmodulin-dependent protein kinase type IV
UniProt Gene Name	CAMK4
UniProt Synonym Gene Names	CAMK; CAMK-GR; CAMKIV; CaMK IV
UniProt Entry Name	KCC4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/camkiv-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/254322>