

Rabbit anti-Mouse IKB beta/NFKBIB Polyclonal Antibody | anti-NFKBIB antibody

Anti-ICK beta/NFKBIB Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA254668	Rabbit	Mouse	Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ICK beta/NFKBIB; Polyclonal Antibody; Anti-ICK beta/NFKBIB Antibody; Rabbit Polyclonal; NFKBIB/IkK-beta Antibody; Rabbit PAb; Antigen Affinity Purified; NFKB inhibitor beta; Anti-I(Kappa)B(beta) Antibody; Anti-I-kappa-B-beta Antibody; Anti-IKappaBbeta Antibody; Anti-IkK Antibody; Anti-IkK-B Antibody; Anti-ICK-beta Antibody; Anti-IkKb Antibody; Anti-NF-kappa-BIB Antibody; anti-NFKBIB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Mouse NFKBIB/IkK-beta
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-NFKBIB antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the C-terminus of the Mouse NFKBIB/IkK-beta
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Mouse NFKBIB/IkK-beta, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of Nfkbib in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti- Nfkbib polyclonal antibody (1:1000) at 4 degree C overnight. Then cells were stained with the Alexa Fluor594-conjugated Goat Anti-rabbit IgG secondary antibody (red). Positive staining was localized to cytoplasm and nucleus.



IHC (Immunohistochemistry)

Immunochemical staining of human Nfkbib in human breast carcinoma with rabbit polyclonal antibody (1:300, formalin-fixed paraffin embedded sections).



ADDITIONAL INFORMATION

NCBI AND UNIPROT INFORMATION

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ikb-beta-nfkbib-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/254668>