

Rabbit CCT3 Polyclonal Antibody | anti-CCT3 antibody

Anti-CCT3 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA254876	Rabbit	Human Predicted to React with: Rat, Cynomolgus (Species predicted to react based on 100% sequence homology)	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CCT3; Polyclonal Antibody; Anti-CCT3 Antibody; Rabbit Polyclonal; CCT3 Antibody; Rabbit PAb; Antigen Affinity Purified; chaperonin containing TCP1; subunit 3 (gamma); Anti-CCT-gamma Antibody; Anti-CCTG Antibody; Anti-PIG48 Antibody; Anti-TCP-1-gamma Antibody; Anti-TRIC5 Antibody; anti-CCT3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human Predicted to React with: Rat, Cynomolgus (Species predicted to react based on 100% sequence homology)
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CCT3
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CCT3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the center region of the Human CCT3
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the center region of the Human CCT3, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human CCT3 in human prostate with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human CCT3 in human colon carcinoma with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-CCT3 rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 61 kDa Observed band size: 61 kDa

NCBI AND UNIPROT INFORMATION

NCBI GI #	58761484
NCBI GeneID	7203
NCBI Accession #	NP_001008800.1
NCBI GenBank Nucleotide #	NM_001008800.2
UniProt Accession #	P49368; Q5SZY1; Q9BR64; A6NE14
Molecular Weight	56,431 Da
NCBI Official Full Name	T-complex protein 1 subunit gamma isoform c
NCBI Official Synonym Full Names	chaperonin containing TCP1 subunit 3
NCBI Official Symbol	CCT3
NCBI Official Synonym Symbols	CCTG; PIG48; TRIC5; CCT-gamma; TCP-1-gamma
NCBI Protein Information	T-complex protein 1 subunit gamma
UniProt Protein Name	T-complex protein 1 subunit gamma
UniProt Gene Name	CCT3
UniProt Synonym Gene Names	CCTG; TRIC5; TCP-1-gamma
UniProt Entry Name	TCPG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cct3-antibody-immunohistochemistry-western-blot-elisa/254876>