

Rabbit anti-Human CSNK2A2 Polyclonal Antibody | anti-CSNK2A2 antibody

Anti-CSNK2A2 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA255291	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, ELISA, Western Blot

PRODUCT SYNONYMS

CSNK2A2; Polyclonal Antibody; Anti-CSNK2A2 Antibody; Rabbit Polyclonal; CSNK2A2 Antibody; Rabbit PAb; Antigen Affinity Purified; casein kinase 2; alpha prime polypeptide; Anti-CK2A2 Antibody; Anti-CSNK2A1 Antibody; anti-CSNK2A2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CSNK2A2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CSNK2A2 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), ELISA, WB (Western Blot)
Immunogen	Recombinant Human CSNK2A2 Protein
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with purified, recombinant Human CSNK2A2 (rh CSNK2A2). CSNK2A2 specific IgG was purified by Human CSNK2A2 affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

CSNK2A2 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate Lane B: 0.5 mg HepG2 Whole Cell Lysate 4 uL anti-CSNK2A2 rabbit polyclonal antibody and 15 ul of 50 % Protein G agarose. Primary antibody: Anti-CSNK2A2 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: DyLight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution. Developed using the Odyssey technique. Performed under reducing conditions. Predicted band size: 41 kDa. Observed band size: 41 kDa.



IF (Immunofluorescence)

Immunofluorescence staining of CSNK2A2 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human CSNK2A2 polyclonal antibody (1:1000) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to cytoplasm and nucleus.



WB (Western Blot)

Anti-CSNK2A2 rabbit polyclonal antibody at 1:500 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: MCF7 Whole Cell Lysate
Lane C: Jurkat Whole Cell Lysate
Lane D: HepG2 Whole Cell Lysate
Lysates/proteins at 30 ug per lane.
Secondary Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.
Developed using the Odyssey technique.
Performed under reducing conditions.
Predicted band size: 41 kDa
Observed band size: 45 kDa
(We are unsure as to the identity of these extra bands.)

ADDITIONAL INFORMATION

Related Product Information for anti-CSNK2A2 antibody	Casein kinase II subunit alpha', also known as CSNK2A2 and CK2A2, is a member of the protein kinase superfamily, Ser/Thr protein kinase family and CK2 subfamily. Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. The alpha and alpha' chains contain the catalytic site. CSNK2A2 is a tetramer composed of an alpha chain, an alpha' and two beta chains. It is also component of a CK2-SPT16-SSRP1 complex composed of SSRP1, SUPT16H, CSNK2A1, CSNK2A2 and CSNK2B, the complex associating following UV irradiation. Protein kinase casein kinase II (Ck2) is a cyclic-AMP and calcium-independent serine-threonine kinase that is composed of two catalytic subunits (alpha and alpha') and two regulatory beta-subunits. Ck2 is not a casein kinase in vivo, but over 1 substrates are known. The highly conserved amino acid sequences of its subunits and their broad expression suggest that Ck2 may have a fundamental role in cell function. Ck2 has been implicated in DNA replication, regulation of basal and inducible transcription, translation and control of metabolism. The Ck2alpha and Ck2alpha' isoforms (products of the genes Csnk2a1 and Csnk2a2, respectively) are highly homologous, the reason for their redundancy and evolutionary conservation is unknown. CSNK2A2 may be a candidate gene for these inherited syndromes.
References	Xu X. et al., 1999, Nat Genet. 23 (1):118-21. Aasland M. et al., 2000, Anim Genet. 31 (2):131-4. Escalier D. et al., 2003, Mol Reprod Dev. 66 (2):190-201. Ackermann K. et al., 2005, Mol Cell Biochem. 274 (1-2):91-101.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503097
NCBI GeneID	1459
NCBI Accession #	NP_001887.1
NCBI GenBank Nucleotide #	NM_001896.2
UniProt Accession #	P19784
Molecular Weight	41kDa
NCBI Official Full Name	casein kinase II subunit alpha'
NCBI Official Synonym Full Names	casein kinase 2 alpha 2
NCBI Official Symbol	CSNK2A2
NCBI Official Synonym Symbols	CK2A2; CSNK2A1; CK2alpha'
NCBI Protein Information	casein kinase II subunit alpha'
UniProt Protein Name	Casein kinase II subunit alpha'
UniProt Gene Name	CSNK2A2
UniProt Synonym Gene Names	CK2A2; CK II alpha'
UniProt Entry Name	CSK22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

