

Rabbit Influenza A H1N1 Hemagglutinin/HA Polyclonal Antibody | anti-HA antibody

Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA Antibody, Rabbit PAb, Antigen Affinity Purified


CATALOG # AAA255310	HOST Rabbit	REACTIVITY H1N1	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	---------------------------	--

PRODUCT SYNONYMS

Influenza A H1N1 Hemagglutinin/HA; Polyclonal Antibody; Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA Antibody; Rabbit PAb; Antigen Affinity Purified; Harvey rat sarcoma viral oncogene homolog; anti-HA antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	H1N1
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA
Purity/Purification	Protein A & Antigen Affinity
Form/Format	0.2um filtered solution in PBS
Concentration	1 mg/ml (varies by lot)
Applicable Applications for anti-HA antibody	ELISA, WB (Western Blot)
Immunogen	Recombinant Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA Protein
Preparation	Produced in rabbits immunized with purified, recombinant Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA (ACP41105.1; Met1-Gln529). Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA specific IgG was purified by Influenza A H1N1 (A/California/04/2009) Hemagglutinin/HA affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.



WB (Western Blot)
Anti-Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA rabbit polyclonal antibody at 1:2000 dilution
Sample: Influenza A H1N1 (A/California/04/2009) Hemagglutinin / HA Recombinant Protein
Lane A: 30ng
Lane B: 10ng
Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution
Performed under reducing conditions.

ADDITIONAL INFORMATION

Related Product Information for anti-HA antibody	The influenza viral Hemagglutinin (HA) protein is a homotrimer with a receptor binding pocket on the globular head of each monomer. HA has at least 18 different antigens. These subtypes are named H1 through H18. HA has two functions. Firstly, it allows the recognition of target vertebrate cells, accomplished through the binding to these cells' sialic acid-containing receptors. Secondly, once bound it facilitates the entry of the viral genome into the target cells by causing the fusion of the host endosomal membrane with the viral membrane. The influenza virus Hemagglutinin (HA) protein is translated in cells as a single protein, HA, or hemagglutinin precursor protein. For viral activation, hemagglutinin precursor protein (HA) must be cleaved by a trypsin-like serine endoprotease at a specific site, normally coded for by a single basic amino acid (usually arginine) between the HA1 and HA2 domains of the protein. After cleavage, the two disulfide-bonded protein domains produce the mature form of the protein subunits as a prerequisite for the conformational change necessary for fusion and hence viral infectivity.
References	White JM, Hoffman LR, Arevalo JH, et al. Attachment and entry of influenza virus into host cells. Pivotal roles of hemagglutinin. In Chiu W, Burnett RM, Garcea RL. Structural Biology of Viruses. 1997 Suzuki Y. Sialobiology of influenza: molecular mechanism of host range variation of influenza viruses. Biol. Pharm. Bull. 2005. Senne DA, Panigrahy B, Kawaoka Y, et al. Survey of the hemagglutinin (HA) cleavage site sequence of H5 and H7 avian influenza viruses: amino acid sequence at the HA cleavage site as a marker of pathogenicity potential. Avian Dis. 1996 Donald J. Benton, Influenza hemagglutinin membrane anchor, PNAS, 2018

NCBI AND UNIPROT INFORMATION

NCBI GI #	227809830
NCBI Official Full Name	hemagglutinin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/influenza-a-h1n1-hemagglutinin-ha-antibody-western-blot-elisa/255310>