

Mouse anti-Human STK33 Monoclonal Antibody | anti-STK33 antibody

STK33 (Serine/Threonine Kinase 33) (HRP)

CATALOG # AAA25560	HOST Mouse	REACTIVITY Human	APPLICATIONS ELISA, Immunoprecipitation, Western Blot
------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

STK33; Monoclonal Antibody; STK33 (Serine/Threonine Kinase 33) (HRP); anti-STK33 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1,k
Clone Number	6G1
Specificity	Recognizes human STK33.
Purity/Purification	Purified by Protein A Affinity Chromatography.
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with horseradish peroxidase (HRP).
Applicable Applications for anti-STK33 antibody	ELISA, IP (Immunoprecipitation), WB (Western Blot)
Application Notes	Applications are based on unconjugated antibody.
Immunogen	Full length recombinant corresponding to aa1-515 from human STK33 (AAH31231) with GST tag. MW of the GST tag alone is 26kD.
Immunogen Sequence	MADSGLDKKSTKCPDCSSASQKDVLCVCSSKTRVPPVLVEMSQTSSIGSAESLISLERKKEKNINRDITSRKDLPSRTSNVERKASQQQWGRGNFTEGH
Conjugate	HRP
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C.

IMAGES



WB (Western Blot)

STK33 monoclonal antibody, Western Blot analysis of STK33 expression in HeLa.



WB (Western Blot)

Western blot analysis of STK33 over-expressed 293 cell line, cotransfected with STK33 Validated Chimera RNAi (Lane 2) or non-transfected control (Lane 1). Blot probed with STK33 monoclonal antibody. GAPDH (36.1kD) used as specificity and loading control.

**IP (Immunoprecipitation)**

Immunoprecipitation of STK33 transfected lysate using STK33 monoclonal antibody and Protein A Magnetic Bead, and immunoblotted with STK33 rabbit polyclonal antibody.

**IF (Immunofluorescence)**

Immunofluorescence of monoclonal antibody to STK33 on HeLa cell. [antibody concentration 25ug/ml].

**WB (Western Blot)**

Western Blot analysis of STK33 expression in transfected 293T cell line by STK33 monoclonal antibody. Lane 1: STK33 transfected lysate (57.9kD). Lane 2: Non-transfected lysate.

**WB (Western Blot)**

Western Blot detection against Immunogen (82.65kD).

ADDITIONAL INFORMATION

Related Product Information for anti-STK33 antibody	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains.
Product Categories/Family for anti-STK33 antibody	Antibodies ; Protein Kinases

NCBI AND UNIPROT INFORMATION

NCBI GI #	22658391
NCBI GeneID	65975
Molecular Weight	50,176 Da
NCBI Official Full Name	Homo sapiens serine/threonine kinase 33, mRNA
NCBI Official Synonym Full Names	serine/threonine kinase 33
NCBI Official Symbol	STK33
NCBI Protein Information	serine/threonine-protein kinase 33

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/stk33-antibody-clone-6g1-immunoprecipitation-western-blot-elisa/25560>