

Rabbit anti-Human TRAF4 Polyclonal Antibody | anti-Traf4 antibody

Anti-TRAF4 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA255864	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

TRAF4; Polyclonal Antibody; Anti-TRAF4 Antibody; Rabbit Polyclonal; TRAF4 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CART1 Antibody; Anti-MLN62 Antibody; Anti-RNF83 Antibody; anti-Traf4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human TRAF4
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-Traf4 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human TRAF4 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human TRAF4 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of TRAF4 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human TRAF4 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



WB (Western Blot)

Anti-TRAF4 rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lane B: Jurkat Whole Cell Lysate Lane C: SKBR-3 Whole Cell Lysate Lane D: 293T Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 54 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-Traf4 antibody	traf4a mRNA is expressed in the developing eye vesicle over the period of optic cup morphogenesis, and Traf4a loss leads to disrupted evagination and elongation of the eye vesicles, and aberrant organization and apico-basal polarity of the eye epithelium. Traf4 acts in a TNF receptor-and Jun N-terminal protein kinase (JNK)-independent manner to fine-tune the assembly of adherens junctions in the invaginating mesodermal cells. Tumor necrosis factor receptor associated factor 4 (Traf4) mRNA expression is upregulated in various breast tumors and tumor cell lines.
Product Categories/Family for anti-Traf4 antibody	IL17 signaling pathway

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/traf4-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/255864>