

Rabbit anti-Human FIGNL1 Polyclonal Antibody | anti-FIGNL1 antibody

Anti-FIGNL1 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA255889	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

FIGNL1; Polyclonal Antibody; Anti-FIGNL1 Antibody; Rabbit Polyclonal; FIGNL1 Antibody; Rabbit PAb; Antigen Affinity Purified; fidgetin-like 1; anti-FIGNL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FIGNL1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FIGNL1 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human FIGNL1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human FIGNL1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IP (Immunoprecipitation)

FIGNL1 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate, 4 μ L anti-FIGNL1 rabbit polyclonal antibody and 60 μ g of Immunomagnetic beads Protein A/G. Primary antibody: Anti-FIGNL1 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 74 kDa. Observed band size: 72 kDa.



IHC (Immunohistochemistry)

Immunochemical staining of human FIGNL1 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-FIGNL1 rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 74 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fign1-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/255889>