

Rabbit anti-Human FKBPL Polyclonal Antibody | anti-FKBPL antibody

Anti-FKBPL Antibody, Rabbit Polyclonal


CATALOG # AAA255892	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---

PRODUCT SYNONYMS


FKBPL; Polyclonal Antibody; Anti-FKBPL Antibody; Rabbit Polyclonal; FKBPL Antibody; Rabbit PAb; Antigen Affinity Purified; FK506 binding protein like; Anti-DIR1 Antibody; Anti-NG7 Antibody; Anti-WISP39 Antibody; anti-FKBPL antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FKBPL
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FKBPL antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human FKBPL fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human FKBPL fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IHC (Immunohistochemistry)
Immunochemical staining of human FKBPL in human corpus uteri with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)
Anti-FKBPL rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lane B: MCF7 Whole Cell Lysate Lane C: HL-60 Whole Cell Lysate Lane D: HeLa Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 38 kDa Observed band size: 44 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fkbpl-antibody-immunohistochemistry-western-blot-elisa/255892>