

Rabbit anti-Human HIBCH Polyclonal Antibody | anti-HIBCH antibody

Anti-HIBCH Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA255951	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HIBCH; Polyclonal Antibody; Anti-HIBCH Antibody; Rabbit Polyclonal; HIBCH Antibody; Rabbit PAb; Antigen Affinity Purified; 3-hydroxyisobutyryl-CoA hydrolase; Anti-HIBYLCOAH Antibody; anti-HIBCH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human HIBCH
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-HIBCH antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human HIBCH fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human HIBCH fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IP (Immunoprecipitation)

HIBCH was immunoprecipitated using: Lane A: 0.5 mg HepG2 Whole Cell Lysate, 4 uL anti-HIBCH rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-HIBCH rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 43 kDa. Observed band size: 43 kDa.




IHC (Immunohistochemistry)

Immunochemical staining of human HIBCH in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human HIBCH in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-HIBCH rabbit polyclonal antibody at 1:500 dilution Lane A: HepG2 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 43 kDa Observed band size: 43 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/hibch-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/255951>