

Rabbit anti-Human COASY Polyclonal Antibody | anti-COASY antibody

Anti-COASY Antibody, Rabbit Polyclonal

CATALOG # AAA255962	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

COASY; Polyclonal Antibody; Anti-COASY Antibody; Rabbit Polyclonal; COASY Antibody; Rabbit PAb; Antigen Affinity Purified; CoA synthase; Anti-DPCK Antibody; Anti-NBIA6 Antibody; Anti-NBP Antibody; Anti-pOV-2 Antibody; Anti-PPAT Antibody; Anti-UKR1 Antibody; anti-COASY antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human COASY
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-COASY antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human COASY fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human COASY fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

COASY was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate, 4 uL anti-COASY rabbit polyclonal antibody and 60 ug of Immnomagnetic beads Protein A/G. Primary antibody: Anti-COASY rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 62 kDa. Observed band size: 62 kDa.



WB (Western Blot)

Anti-COASY rabbit polyclonal antibody at 1:500 dilution. Lane A: HepG2 Whole Cell Lysate. Lane B: HeLa Whole Cell Lysate. Lane C: A431 Whole Cell Lysate. Lane D: HEK293 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 62 kDa. Observed band size: 62 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.