

Rabbit anti-Human CAPZB Polyclonal Antibody | anti-CAPZB antibody

Anti-CAPZB Antibody, Rabbit Polyclonal



CATALOG # AAA255994	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

CAPZB; Polyclonal Antibody; Anti-CAPZB Antibody; Rabbit Polyclonal; CAPZB Antibody; Rabbit PAb; Antigen Affinity Purified; capping protein (actin filament) muscle Z-line; beta; Anti-CAPB Antibody; Anti-CAPPB Antibody; Anti-CAPZ Antibody; anti-CAPZB antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CAPZB
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CAPZB antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human CAPZB fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human CAPZB fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)
CAPZB was immunoprecipitated using: Lane A: 0.5 mg A431 Whole Cell Lysate 4 uL anti-CAPZB rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CAPZB rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 31 kDa. Observed band size: 31 kDa.

WB (Western Blot)
Anti-CAPZB rabbit polyclonal antibody at 1:500 dilution. Lane A: HeLa Whole Cell Lysate. Lane B: 293T Whole Cell Lysate. Lane C: Jurkat Whole Cell Lysate. Lane D: A431 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 31 kDa. Observed band size: 31 kDa. (We are unsure as to the identity of these extra bands.)

ADDITIONAL INFORMATION

Product Categories/Family for anti-CAPZB antibody	Actin Dynamics Signaling Pathway
---	----------------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/capzb-antibody-immunoprecipitation-western-blot-elisa/255994>