

Rabbit anti-Human MAGOH Polyclonal Antibody | anti-MAGOH antibody

Anti-MAGOH Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256086	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAGOH; Polyclonal Antibody; Anti-MAGOH Antibody; Rabbit Polyclonal; MAGOH Antibody; Rabbit PAb; Antigen Affinity Purified; mago-nashi homolog; proliferation-associated (Drosophila); Anti-MAGOH1 Antibody; Anti-MAGOHA Antibody; anti-MAGOH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human MAGOH
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-MAGOH antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human MAGOH fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human MAGOH fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

MAGOH was immunoprecipitated using: Lane A: 0.5 mg K562 Whole Cell Lysate 4 uL anti-MAGOH rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-MAGOH rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 17 kDa. Observed band size: 17 kDa.




IF (Immunofluorescence)

Immunofluorescence staining of MAGOH in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human MAGOH polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Nucleus and Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human MAGOH in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-MAGOH rabbit polyclonal antibody at 1:2000 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: Jurkat Whole Cell Lysate
Lane C: A431 Whole Cell Lysate
Lane D: K562 Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 17 kDa. Observed band size: 17 kDa (We are unsure as to the identity of these extra bands).

ADDITIONAL INFORMATION

Related Product Information for anti-MAGOH antibody

MAGOH (Mago Homolog, Exon Junction Complex Subunit) is a Protein Coding gene. MAGOH is the human homolog of *Drosophila mago nashi*, a protein that is required for normal germline development in the *Drosophila* embryo. In mammals, mRNA expression is not limited to the germline, but is expressed ubiquitously in adult tissues and can be induced by serum stimulation of quiescent fibroblasts. The exon junction complex (EJCs) are deposited on messenger RNAs (mRNAs) during splicing and their core consists of eIF4A3, MLN51, Y14, and MAGOH. Both MAGOH proteins interact with other EJC components, incorporate into mRNA-bound EJCs, and activate nonsense-mediated decay. Diseases associated with MAGOH include Metaphyseal Chondrodysplasia, Schmid Type, and Primary Autosomal Recessive Microcephaly.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/magoh-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/256086>