

Rabbit anti-Human Mannan Binding Lectin/MBL2 Polyclonal Antibody | anti-MBL2 antibody

Anti-Mannan Binding Lectin/MBL2 Antibody, Rabbit Polyclonal

CATALOG # AAA256098	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Mannan Binding Lectin/MBL2; Polyclonal Antibody; Anti-Mannan Binding Lectin/MBL2 Antibody; Rabbit Polyclonal; Mannan Binding Lectin/MBL2 Antibody; Rabbit PAb; Antigen Affinity Purified; mannose-binding lectin (protein C) 2; soluble; Anti-COLEC1 Antibody; Anti-HSMBPC Antibody; Anti-MBL Antibody; Anti-MBL2D Antibody; Anti-MBP Antibody; Anti-MBP-C Antibody; Anti-MBP1 Antibody; Anti-MBPD Antibody; anti-MBL2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Mannan Binding Lectin/MBL2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-MBL2 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human Mannan Binding Lectin/MBL2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Mannan Binding Lectin/MBL2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunochemical staining of human MBL2 in human kidney with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.


IHC (Immunohistochemistry)

Immunochemical staining of human MBL2 in human liver with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-MBL2 antibody	MBL (mannose-binding lectin) is primarily a liver-derived collagen-like serum protein, which binds sugar structures on micro-organisms and dying host cells and is one of the four known mediators that initiate activation of the complement system via the lectin pathway. MBL and the ficolins (Ficolin-1, Ficolin-2, and Ficolin-3) are soluble collagen-like proteins that are involved in innate immune defense. They bind sugar structures or acetylated compounds present on microorganisms and dying host cells and they initiate activation of the lectin complement pathway in varying degrees. MBL2 encodes the mannose-binding lectin, which is a key player in the innate immune system and has recently been found to play a role in the development of type 1 diabetes and gestational diabetes mellitus. Common variant alleles situated both in the promoter and structural regions of the MBL2 gene influence the stability and the serum concentration of the protein. Several polymorphisms in the promoter and structural regions of MBL2 adversely affect the plasma concentration and the oligomeric state of MBL. The possession of mutant alleles has been linked to disease outcomes for a variety of bacterial and viral infections. Mutant MBL2 haplotypes have been linked to disease progression and response to therapy in HCV infection.
Product Categories/Family for anti-MBL2 antibody	Complement Activation Pathways
References	Garred P, et al. (2006) Mannose-binding lectin and its genetic variants. Genes Immun. 7(2): 85-94.Brown KS, et al. (2007) Mannan binding lectin and viral hepatitis. Immunol Lett. 108(1): 34-44.Garred P. (2008) Mannose-binding lectin genetics: from A to Z. Biochem Soc Trans. 36(Pt 6): 1461-6.Garred P, et al. (2009) MBL2, FCN1, FCN2 and FCN3-The genes behind the initiation of the lectin pathway of complement. Mol Immunol. 46(14): 2737-44.Muller YL, et al. (2010) Functional Variants in MBL2 Are Associated With Type 2 Diabetes and Pre-Diabetes Traits in Pima Indians and the Old Order Amish. Diabetes. 59(8): 2080-5.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mannan-binding-lectin-mbl2-antibody-immunohistochemistry-elisa/256098>