

Rabbit anti-Human Fractalkine Polyclonal Antibody | anti-CX3CL1 antibody

Anti-Fractalkine Antibody, Rabbit Polyclonal

CATALOG # AAA256104	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Fractalkine; Polyclonal Antibody; Anti-Fractalkine Antibody; Rabbit Polyclonal; Fractalkine/CX3CL1 Antibody; Rabbit PAb; Antigen Affinity Purified; chemokine (C-X3-C motif) ligand 1; Anti-ABCD-3 Antibody; Anti-C3Xkine Antibody; Anti-CXC3 Antibody; Anti-CXC3C Antibody; Anti-fractalkine Antibody; Anti-neurotactin Antibody; Anti-NTN Antibody; Anti-NTT Antibody; Anti-SCYD1 Antibody; anti-CX3CL1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Fractalkine/CX3CL1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CX3CL1 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human Fractalkine/CX3CL1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Fractalkine/CX3CL1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunochemical staining of human CX3CL1 in human skeletal muscle with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.


IHC (Immunohistochemistry)

Immunochemical staining of human CX3CL1 in human kidney with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-CX3CL1 antibody	Fractalkine or Chemokine (C-X3-C motif) ligand 1 (CX3CL1) is a member of the CX3C chemokine family. Fractalkine/CX3CL1 is a unique chemokine that functions not only as a chemoattractant but also as an adhesion molecule and is expressed on endothelial cells activated by proinflammatory cytokines, such as interferon-gamma and tumor necrosis factor-alpha. Fractalkine/CX3CL1 is expressed in a membrane-bound form on activated endothelial cells and mediates attachment and firm adhesion of T cells, monocytes and NK cells. Fractalkine/CX3CL1 is associated with dendritic cells (DC) in epidermis and lymphoid organs. The fractalkine receptor, CX3CR1, is expressed on cytotoxic effector lymphocytes, including natural killer (NK) cells and cytotoxic T lymphocytes, which contain high levels of intracellular perforin and granzyme B, and on macrophages. Soluble fractalkine causes migration of NK cells, cytotoxic T lymphocytes, and macrophages, whereas the membrane-bound form captures and enhances the subsequent migration of these cells in response to secondary stimulation with other chemokines.
Product Categories/Family for anti-CX3CL1 antibody	Neuroinflammation
References	Imai T, et al. (1997) Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. Cell. 91(4): 521-30. Papadopoulos EJ, et al. (1999) Fractalkine, a CX3C chemokine, is expressed by dendritic cells and is up-regulated upon dendritic cell maturation. Eur J Immunol. 29 (8): 2551-9. Umehara H, et al. (2004) Fractalkine in vascular biology: from basic research to clinical disease". Arterioscler. Thromb Vasc Biol. 24 (1): 34-40.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fractalkine-antibody-immunohistochemistry-elisa/256104>