

Rabbit anti-Human IFT57/HIPPI Polyclonal Antibody | anti-IFT57 antibody

Anti-IFT57/HIPPI Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256115	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

IFT57/HIPPI; Polyclonal Antibody; Anti-IFT57/HIPPI Antibody; Rabbit Polyclonal; IFT57/HIPPI Antibody; Rabbit PAb; Antigen Affinity Purified; intraflagellar transport 57; Anti-ESRRBL1 Antibody; Anti-HIPPI Antibody; Anti-MHS4R2 Antibody; anti-IFT57 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human IFT57/HIPPI
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-IFT57 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human IFT57/HIPPI fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human IFT57/HIPPI fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

IFT57 was immunoprecipitated using: Lane A: 0.5 mg Raji Whole Cell Lysate 4 uL anti-IFT57 rabbit polyclonal antibody and 60 ug of Immnomagnetic beads Protein A/G. Primary antibody: Anti-IFT57 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 49 kDa. Observed band size: 55 kDa.



WB (Western Blot)

Anti-IFT57 rabbit polyclonal antibody at 1:500 dilution. Lane A: Raji Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 49 kDa. Observed band size: 49 kDa. (We are unsure as to the identity of these extra bands.)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.