

Rabbit anti-Human CLNS1A Polyclonal Antibody | anti-CLNS1A antibody

Anti-CLNS1A Antibody, Rabbit Polyclonal

CATALOG # AAA256138	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CLNS1A; Polyclonal Antibody; Anti-CLNS1A Antibody; Rabbit Polyclonal; CLNS1A Antibody; Rabbit PAb; Antigen Affinity Purified; chloride channel; nucleotide-sensitive; 1A; Anti-CLCI Antibody; Anti-CLNS1B Antibody; Anti-ICln Antibody; anti-CLNS1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CLNS1A
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CLNS1A antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human CLNS1A fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human CLNS1A fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

CLNS1A was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate 4 uL anti-CLNS1A rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CLNS1A rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 26 kDa. Observed band size: 40 kDa.



WB (Western Blot)

Anti-CLNS1A rabbit polyclonal antibody at 1:500 dilution. Lane A: Jurkat Whole Cell Lysate. Lane B: HeLa Whole Cell Lysate. Lane C: K562 Whole Cell Lysate. Lane D: HepG2 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 26 kDa. Observed band size: 38 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.