

Rabbit anti-Human CD2AP Polyclonal Antibody | anti-CD2AP antibody

Anti-CD2AP Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256203	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


CD2AP; Polyclonal Antibody; Anti-CD2AP Antibody; Rabbit Polyclonal; CD2AP Antibody; Rabbit PAb; Antigen Affinity Purified; CD2-associated protein; Anti-CMS Antibody; anti-CD2AP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CD2AP
Purity/Purification	Protein A & Antigen Affinity
Form/Format	PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CD2AP antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human CD2AP fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human CD2AP fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.




IP (Immunoprecipitation)
CD2AP was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate 4 uL anti-CD2AP rabbit polyclonal antibody and 60 ug of Immnomagnetic beads Protein A/G. Primary antibody: Anti-CD2AP rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 71 kDa. Observed band size: 80 kDa.



IHC (Immunohistochemistry)
Immunohistochemical staining of human CD2AP in human pancreas with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)
Immunohistochemical staining of human CD2AP in human placenta with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)
Anti-CD2AP rabbit polyclonal antibody at 1:2000 dilution. Lane A: HeLa Whole Cell Lysate. Lane B: 293T Whole Cell Lysate. Lane C: Jurkat Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 71 kDa. Observed band size: 84 kDa.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD2AP antibody	Adaptor Proteins Apoptosis Adaptor Proteins
---	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.