

Rabbit anti-Human ACADVL Polyclonal Antibody | anti-ACADVL antibody

Anti-ACADVL Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256272	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry

PRODUCT SYNONYMS

ACADVL; Polyclonal Antibody; Anti-ACADVL Antibody; Rabbit Polyclonal; ACADVL Antibody; Rabbit PAb; Antigen Affinity Purified; acyl-CoA dehydrogenase; very long chain; Anti-ACAD6 Antibody; Anti-LCACD Antibody; Anti-VLCAD Antibody; anti-ACADVL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human ACADVL
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ACADVL antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human ACADVL fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human ACADVL fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.




IF (Immunofluorescence)

Immunofluorescence staining of ACADVL in A549 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human ACADVL polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human ACADVL in human adrenal gland with rabbit polyclonal antibody at 1:10000 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human ACADVL in human duodenum with rabbit polyclonal antibody at 1:10000 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/acadvl-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/256272>