

Rabbit anti-Human RhoGDI/ARHGDI Polyclonal Antibody | anti-ARHGDI antibody

Anti-RhoGDI/ARHGDI Antibody, Rabbit Polyclonal



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256492	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

RhoGDI/ARHGDI; Polyclonal Antibody; Anti-RhoGDI/ARHGDI Antibody; Rabbit Polyclonal; RhoGDI/ARHGDI Antibody; Rabbit PAb; Antigen Affinity Purified; Rho GDP dissociation inhibitor (GDI) alpha; Anti-GDIA1 Antibody; Anti-HEL-S-47e Antibody; Anti-NPHS8 Antibody; Anti-RHOGDI Antibody; Anti-RHOGDI-1 Antibody; anti-ARHGDI antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human RhoGDI/ARHGDI
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ARHGDI antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human RhoGDI/ARHGDI fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human RhoGDI/ARHGDI fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.




IP (Immunoprecipitation)

ARHGDIA was immunoprecipitated using: Lane A: 0.5 mg HL-60 Whole Cell Lysate 4 uL anti-ARHGDIA rabbit polyclonal antibody and 60 ug of Immnomagnetic beads Protein A/G. Primary antibody: Anti-ARHGDIA rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 23 kDa Observed band size :25 kDa

IHC (Immunohiostchemistry)

Immunochemical staining of human ARHGDIA in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-ARHGDIA rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lane B: HeLa Whole Cell Lysate Lane C: HL-60 Whole Cell Lysate Lane D: U-251 MG Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 23 kDa Observed band size: 26 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.