

Rabbit anti-Human HMGCL Polyclonal Antibody | anti-HMGCL antibody

Anti-HMGCL Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256658	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry

PRODUCT SYNONYMS

HMGCL; Polyclonal Antibody; Anti-HMGCL Antibody; Rabbit Polyclonal; HMGCL Antibody; Rabbit PAb; Antigen Affinity Purified; 3-hydroxymethyl-3-methylglutaryl-CoA lyase; Anti-HL Antibody; anti-HMGCL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human HMGCL
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-HMGCL antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human HMGCL fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human HMGCL fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.




IF (Immunofluorescence)

Immunofluorescence staining of HMGCL in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human HMGCL polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human HMGCL in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human HMGCL in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/hmgcl-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/256658>