

Rabbit anti-Human DTYMK Polyclonal Antibody | anti-DTYMK antibody

Anti-DTYMK Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256702	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

DTYMK; Polyclonal Antibody; Anti-DTYMK Antibody; Rabbit Polyclonal; DTYMK Antibody; Rabbit PAb; Antigen Affinity Purified; deoxythymidylate kinase; Anti-CDC8 Antibody; Anti-PP3731 Antibody; Anti-TMPK Antibody; Anti-TYMK Antibody; anti-DTYMK antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human DTYMK
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-DTYMK antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human DTYMK fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human DTYMK fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.




IP (Immunoprecipitation)

DTYMK was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate 4 uL anti-DTYMK rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-DTYMK rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 23 kDa. Observed band size: 26 kDa.




IHC (Immunohistochemistry)

Immunochemical staining of human DTYMK in human stomach with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human DTYMK in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-DTYMK rabbit polyclonal antibody at 1:500 dilution. Lane A: HeLa Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 23 kDa. Observed band size: 26 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dtymk-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/256702>