

Rabbit anti-Human REXO1 Polyclonal Antibody | anti-REXO1 antibody

Anti-REXO1 Antibody, Rabbit Polyclonal

CATALOG # AAA256724	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

REXO1; Polyclonal Antibody; Anti-REXO1 Antibody; Rabbit Polyclonal; REXO1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-EloA-BP1 Antibody; Anti-ELOABP1 Antibody; Anti-REX1 Antibody; Anti-TCEB3BP1 Antibody; anti-REXO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human REXO1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-REXO1 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human REXO1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human REXO1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human REXO1 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human REXO1 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-REXO1 antibody	The Elongin complex stimulates the rate of transcription elongation by RNA polymerase II by suppressing the transient pausing of the polymerase at many sites along the DNA template. Elongin is composed of a transcriptionally active A subunit, and two positive regulatory B and C subunits. Although the NH(2)-terminal approximately 120 amino acid region of Elongin A is dispensable for its transcriptional activity in vitro, it shares significant sequence similarity with the NH(2)-terminus of other class of transcription factors SII and CRSP70. Elongin A-binding protein 1 (EloA-BP1), a human exonuclease domain-containing protein, is composed of 1221 amino acids and its mRNA is ubiquitously expressed.
References	Tamura K, et al. (2003) Identification of elo-a-bp1, a novel elongin a binding protein with an exonuclease homology domain. Biochem Biophys Res Commun 309 (1): 189-195.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rexo1-antibody-immunohistochemistry-elisa/256724>