

Rabbit anti-Human LYPD2 Polyclonal Antibody | anti-LYPD2 antibody

Anti-LYPD2 Antibody, Rabbit Polyclonal

CATALOG # AAA256786	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

LYPD2; Polyclonal Antibody; Anti-LYPD2 Antibody; Rabbit Polyclonal; LYPD2 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-LYPDC2 Antibody; Anti-UNQ430 Antibody; Anti-UNQ430/PRO788 Antibody; anti-LYPD2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human LYPD2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-LYPD2 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human LYPD2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human LYPD2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunochemical staining of human LYPD2 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.


IHC (Immunohistochemistry)

Immunochemical staining of human LYPD2 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-LYPD2 antibody	LYPD2 (LY6/PLAUR Domain Containing 2, Also known as LYPDC2; UNQ430) is located on human chromosome 8q24.3. The LYPD2 protein is biasedly expressed in esophagus, stomach and other tissue. Structurally, the encoded LYPD2 protein is reported to be 13115 Da in mass. 68 organisms have orthologs with human gene LYPD2. LYPD2 is associated with some diseases, such as Mal De Meleda. Among its related pathways are Metabolism of proteins and Post-translational modification-synthesis of GPI-anchored proteins. An important paralog of this gene is SLURP1.
References	Su PH, et al. (2017) Methyloomics of nitroxidative stress on precancerous cells reveals DNA methylation alteration at the transition from in situ to invasive cervical cancer. Oncotarget 8 (39): 65281-65291.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/lypd2-antibody-immunohistochemistry-elisa/256786>