

Rabbit anti-Human A33 Polyclonal Antibody | anti-GPA33 antibody

Anti-A33 Antibody, Rabbit Polyclonal

CATALOG # AAA256912	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

A33; Polyclonal Antibody; Anti-A33 Antibody; Rabbit Polyclonal; A33/GPA33 Antibody; Rabbit PAb; Antigen Affinity Purified; glycoprotein A33 (transmembrane); anti-GPA33 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human A33/GPA33
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-GPA33 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human A33/GPA33 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human A33/GPA33 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical staining of human A33 in human small intestine with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.


IHC (Immunohistochemistry)

Immunohistochemical staining of human A33 in human rectum with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-GPA33 antibody	Cell surface A33 antigen, also known as glycoprotein A33, is a single-pass type I membrane protein that is expressed in the normal gastrointestinal epithelium and 95% of colon cancers. GPA33 contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. The open reading frame encodes a 319-amino acid polypeptide having a putative secretory signal sequence and 3 potential glycosylation sites. The predicted mature protein has a 213-amino acid extracellular region, a single transmembrane domain, and a 62-amino acid intracellular tail. Intracellular traffic and recycling to the cell surface appear to play a major role in GPA33 function and to have an influence on its surface density superseding translational regulation. GPA33 has become a promising target of immunologic therapy strategies, but its biologic function and potential role in tumorigenesis are unknown. EpCAM protein and GPA33 mRNA expressions are specific and sensitive markers of Barrett's metaplasia (BM). GPA33 may also play a role in cell-cell recognition and signaling.
Product Categories/Family for anti-GPA33 antibody	Cancer Drug Targets
References	Heath J.K., et al., 1997, Proc. Natl. Acad. Sci. USA. 94:469-74. Ritter G., et al., 1997, Biochem. Biophys. Res. Commun. 236:682-6. Frey, D. et al., 2008, Cancer Biother Radiopharm 23 (1):65-73. Rageul, J. et al., 2009, Int J Cancer 125 (12):2802-9.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/a33-antibody-immunohistochemistry-elisa/256912>