

Rabbit anti-Mouse LDLR/LDL Receptor Monoclonal Antibody | anti-LDLR antibody

Recombinant Anti-LDLR/LDL Receptor Antibody (FITC), Rabbit Monoclonal


CATALOG # AAA257131	HOST Rabbit	REACTIVITY Mouse	APPLICATIONS Flow Cytometry, Functional Assay
-------------------------------	-----------------------	----------------------------	---

PRODUCT SYNONYMS

LDLR/LDL Receptor; Monoclonal Antibody; Recombinant Anti-LDLR/LDL Receptor Antibody (FITC); Rabbit Monoclonal; LDLR/LDL Receptor Antibody (FITC); Rabbit MAb; low density lipoprotein receptor; Anti-Hlb301 Antibody; anti-LDLR antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	032
Specificity	Mouse LDLR/LDL Receptor
Purity/Purification	Protein A
Form/Format	Liquid; Aqueous solution containing 0.5% BSA and 0.09% sodium azide
Concentration	10ul/Test, 0.1mg/ml (varies by lot)
Applicable Applications for anti-LDLR antibody	FCM/FACS (Flow Cytometry)
Antibody Type	Recombinant Antibody
Immunogen	Recombinant Mouse LDLR/LDL Receptor protein
Conjugation	FITC
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Mouse LDLR/LDL Receptor (rM LDLR/LDL Receptor) and conjugated with FITC under optimum conditions, the unreacted FITC was removed.
Preparation and Storage	This antibody can be stored at 2-8 degree C for twelve months without detectable loss of activity. Protected from prolonged exposure to light. Do not freeze! Sodium azide is toxic to cells and should be disposed of properly. Flush with large volumes of water during disposal.



Application Data
Profile of anti-LDLR reactivity on Raw264.7 cells analyzed by flow cytometry. Cells should be Fc-blocked by treatment with Mouse BD Fc Block[®] purified anti-CD16/CD32 mAb 2.4G2 prior to staining, washed, then stained with FITC Rabbit anti-LDLR.

ADDITIONAL INFORMATION

Related Product Information for anti-LDLR antibody	LDL Receptor, also known as LDLR, is a mosaic protein that belongs to the Low-density lipoprotein receptor gene family. The low-density lipoprotein receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. LDL Receptor consists of 840 amino acids (after removal of signal peptide) and mediates the endocytosis of cholesterol-rich LDL. Low-density lipoprotein (LDL) is normally bound at the cell membrane and taken into the cell ending up in lysosomes where the protein is degraded and the cholesterol is made available for repression of microsomal enzyme 3-hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase, the rate-limiting step in cholesterol synthesis. At the same time, a reciprocal stimulation of cholesterol ester synthesis takes place. LDL Receptor is a cell-surface receptor that recognizes the apoprotein B100 which is embedded in the phospholipid outer layer of LDL particles. The receptor also recognizes the apoE protein found in chylomicron remnants and VLDL remnants.
References	Yamamoto T, et al. (1984) The human LDL receptor: a cysteine-rich protein with multiple Alu sequences in its mRNA. Cell. 39(1): 27-38.Mao B, et al. (2001) LDL-receptor-related protein 6 is a receptor for Dickkopf proteins. Nature. 411(6835): 321-5.Pinson KI, et al. (2000) An LDL-receptor-related protein mediates Wnt signalling in mice. Nature. 407(6803): 535-8.

NCBI AND UNIPROT INFORMATION

NCBI GI #	113195700
NCBI GeneID	16835
NCBI Accession #	NP_034830.2
NCBI GenBank Nucleotide #	NM_010700.3
UniProt Accession #	P35951
NCBI Official Full Name	low-density lipoprotein receptor isoform 1
NCBI Official Synonym Full Names	low density lipoprotein receptor
NCBI Official Symbol	Ldlr
NCBI Official Synonym Symbols	Hlb301
NCBI Protein Information	low-density lipoprotein receptor; LDL receptor
UniProt Protein Name	Low-density lipoprotein receptor
UniProt Gene Name	Ldlr
UniProt Synonym Gene Names	LDL receptor

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ldlr-ldl-receptor-antibody-clone-032-flow-cytometry-elisa/257131>