

Rabbit anti-Mouse ICAM-1 Monoclonal Antibody | anti-ICAM1 antibody

Recombinant Anti-ICAM-1 Antibody, Rabbit Monoclonal

CATALOG # AAA257157	HOST Rabbit	REACTIVITY Mouse	APPLICATIONS Immunohistochemistry, ELISA, ELISA
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

ICAM-1; Monoclonal Antibody; Recombinant Anti-ICAM-1 Antibody; Rabbit Monoclonal; ICAM-1/CD54 Antibody; Rabbit MAb; intercellular adhesion molecule 1; Anti-CD54 Antibody; Anti-Icam-1 Antibody; Anti-Ly-47 Antibody; Anti-MALA-2 Antibody; anti-ICAM1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	020
Specificity	Mouse CD54/ICAM1. No cross-reactivity in ELISA with Human ICAM1, Human CD31, Human ICAM3, Human ICAM2, Mouse VCAM1
Purity/Purification	Protein A
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-ICAM1 antibody	IHC (Immunohistochemistry), ELISA, ELISA
Antibody Type	Recombinant Antibody
Immunogen	Recombinant Mouse CD54/ICAM1 protein
Conjugation	Unconjugated
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Mouse CD54/ICAM1 (rM CD54/ICAM1).
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of mouse ICAM1 in mouse brain with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections). The image showing positive staining of choroid plexus.



IHC (Immunohistochemistry)

Immunochemical staining of mouse ICAM1 in mouse kidney with rabbit monoclonal antibody (1:200, formalin-fixed paraffin embedded sections). Positive staining was localized to blood vessel.

ADDITIONAL INFORMATION

Related Product Information for anti-ICAM1 antibody	Intercellular adhesion molecule-1 (ICAM-1, or CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and animal models of atherosclerosis. Additionally, ICAM-1 has been implicated in the progression of autoimmune diseases. ICAM-1 is a ligand for LFA-1(integrin). When activated, leukocytes bind to endothelial cells via ICAM-1/LFA-1 interaction and then transmigrate into tissues. Presence with heavy glycosylation and other structural characteristics, ICAM-1 possesses binding sites for some immune-associated ligands and serves as the binding site for entry of the major group of human Rhinovirus (HRV) into various cell types. ICAM-1 also becomes known for its affinity for Plasmodium falciparum-infected erythrocytes (PFIE), providing more of a role in infectious disease. Previous studies have shown that ICAM-1 is involved in inflammatory reactions and that a defect in ICAM-1 gene inhibits allergic contact hypersensitivity.
Product Categories/Family for anti-ICAM1 antibody	Neuroinflammation
References	Xu H, et al. (2001) The role of ICAM-1 molecule in the migration of Langerhans cells in the skin and regional lymph node. Eur J Immunol. 31(10): 3085-93.Terol MJ, et al. (2003) Soluble intercellular adhesion molecule-1 (s-ICAM-1/s-CD54) in diffuse large B-cell lymphoma: association with clinical characteristics and outcome. Ann Oncol. 14(3): 467-74.Mendez MP, et al. (2006) Shedding of soluble ICAM-1 into the alveolar space in murine models of acute lung injury. Am J Physiol Lung Cell Mol Physiol. 290(5): L962-70.Lawson C, et al. (2009) ICAM-1 signaling in endothelial cells. Pharmacol Rep. 61(1): 22-32.

NCBI AND UNIPROT INFORMATION

NCBI GI #	21389311
NCBI GeneID	15894
NCBI Accession #	NP_034623.1
NCBI GenBank Nucleotide #	NM_010493.2
UniProt Accession #	P13597; Q61828
Molecular Weight	58,753 Da
NCBI Official Full Name	intercellular adhesion molecule 1
NCBI Official Synonym Full Names	intercellular adhesion molecule 1
NCBI Official Symbol	Icam1
NCBI Official Synonym Symbols	CD54; Ly-47; Icam-1; MALA-2
NCBI Protein Information	intercellular adhesion molecule 1; myD10
UniProt Protein Name	Intercellular adhesion molecule 1
UniProt Gene Name	Icam1
UniProt Synonym Gene Names	Icam-1; ICAM-1
UniProt Entry Name	ICAM1_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.