

VEGFD recombinant protein

VEGFD Protein, Human, Recombinant (His Tag)

CATALOG #
AAA257426



HOST
HEK293 Cells

PRODUCT SYNONYMS

VEGFD; VEGFD Protein; Human; Recombinant (His Tag); Human VEGF-D/VEGFD/FIGF Protein (His Tag); c-fos induced growth factor (vascular endothelial growth factor D); VEGF-D Protein; VEGFD recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>95 % as determined by SDS-PAGE.
Form/Format	Lyophilized from sterile PBS, pH7.4 Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Sequence Positions	120aa
Sequence	Phe93-Ser201
Species	Human
Tag	C-His
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Phe93
Protein Construction	A DNA sequence encoding the mature form of human VEGFD (NP_004460.1) corresponding to amino acid (Phe 93-Ser 201) was expressed with a C-terminal polyhistidine tag
Activity	Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC). The ED50 for this effect is 0.3-1.6ug/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. Shipping: They are shipped out with blue ice pack



Application Data

Application Data
Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC).
The ED50 for this effect is 0.3-1.6 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for VEGFD recombinant protein	Vascular endothelial growth factor D (VEGF-D), also known as C-fos induced growth factor (FIGF), belongs to the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family. FIGF protein is active in angiogenesis, lymphangiogenesis, and endothelial cell growth. FIGF protein is secreted as a non-covalent homodimer in an antiparallel fashion. Human FIGF protein is expressed in adult lung, heart, muscle, and small intestine, and is most abundantly expressed in fetal lungs and skin. FIGF protein is structurally and functionally similar to VEGF-C. Therefore, FIGF protein binds and activates VEGFR-2 (Flk1) and VEGFR-3 (Flt4) receptors, and may particularly be involved in cancers, such as breast cancer, epithelial ovarian carcinoma and so on.
References	Avantaggiato V, et al. (1998) Embryonic expression pattern of the murine figf gene, a growth factor belonging to platelet-derived growth factor/vascular endothelial growth factor family. Mech Dev. 73(2):221-4. Rocchigiani M, et al. (1998) Human FIGF: cloning, gene structure, and mapping to chromosome Xp22.1 between the PIGA and the GRPR genes. Genomics 47(2):207-16. Karpanen T, et al. (2008) VEGF-D: a modifier of embryonic lymphangiogenesis. Blood. 112(5): 1547-8.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758378
NCBI GeneID	2277
NCBI Accession #	NP_004460.1
NCBI GenBank Nucleotide #	NM_004469.4
UniProt Accession #	O43915; B2R7Z3
Molecular Weight	40,444 Da
NCBI Official Full Name	vascular endothelial growth factor D preproprotein
NCBI Official Synonym Full Names	c-fos induced growth factor (vascular endothelial growth factor D)
NCBI Official Symbol	FIGF
NCBI Official Synonym Symbols	VEGFD; VEGF-D
NCBI Protein Information	vascular endothelial growth factor D
UniProt Protein Name	Vascular endothelial growth factor D
UniProt Gene Name	FIGF
UniProt Synonym Gene Names	VEGFD; VEGF-D; FIGF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/figf-recombinant-protein-hek293-cells/257426>