

Rabbit anti-Human FANCC Polyclonal Antibody | anti-FANCC antibody

Anti-FANCC Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257498	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry


PRODUCT SYNONYMS

FANCC; Polyclonal Antibody; Anti-FANCC Antibody; Rabbit Polyclonal; FANCC Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-FA3 Antibody; Anti-FAC Antibody; Anti-FACC Antibody; Fanconi anemia; complementation group C; anti-FANCC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FANCC
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. 0.2um filtered solution in PBS
Applicable Applications for anti-FANCC antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human FANCC
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human FANCC, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human FANCC in human brain with rabbit polyclonal antibody (1:2000, formalin-fixed paraffin embedded sections).



IHC (Immunohistochemistry)

Immunochemical staining of human FANCC in human pancreas with rabbit polyclonal antibody (1:2000, formalin-fixed paraffin embedded sections).



IF (Immunofluorescence)

Immunofluorescence staining of FANCC in HeLa cells. Cells were fixed with 4% PFA, blocked with 10% serum. Then incubated with rabbit anti-human FANCC polyclonal antibody (1:1000) (Figure A), incubated with rabbit anti-human FANCC polyclonal antibody and antigen (Figure B) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to cytoplasm and nucleus.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fancc-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/257498>