

Rabbit anti-Human DNAJB11/ERdj3 Polyclonal Antibody | anti-DNAJB11 antibody

Anti-DNAJB11/ERdj3 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257552	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DNAJB11/ERdj3; Polyclonal Antibody; Anti-DNAJB11/ERdj3 Antibody; Rabbit Polyclonal; DNAJB11/ERdj3 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-ABBP-2 Antibody; Anti-ABBP2 Antibody; Anti-Dj-9 Antibody; Anti-DJ9 Antibody; Anti-EDJ Antibody; Anti-ERdj3 Antibody; Anti-ERj3 Antibody; Anti-ERj3p Antibody; Anti-PRO1080 Antibody; Anti-UNQ537 Antibody; DnaJ (Hsp40) homolog; subfamily B; member 11; anti-DNAJB11 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human DNAJB11/ERdj3
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-DNAJB11 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human DNAJB11/ERdj3 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human DNAJB11/ERdj3 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES



IP (Immunoprecipitation)

DNAJB11 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate 4 uL anti-DNAJB11 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-DNAJB11 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 40 kDa. Observed band size: 42 kDa.



IHC (Immunohistochemistry)

Immunohistochemical staining of human DNAJB11 in human pancreas with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-DNAJB11/ERdj3 rabbit polyclonal antibody at 1:500 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: COLO205 Whole Cell Lysate
Lane C: MCF-7 Whole Cell Lysate
Lysates/proteins at 30 ug per lane.
Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.
Developed using the ECL technique.
Performed under reducing conditions.
Predicted band size: 40 kDa
Observed band size: 42 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/dnajb11-erdj3-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/257552>