

Rabbit anti-Human Cytokeratin 20/KRT20 Polyclonal Antibody | anti-KRT20 antibody

Anti-Cytokeratin 20/KRT20 Antibody, Rabbit Polyclonal


CATALOG # AAA257558	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

Cytokeratin 20/KRT20; Polyclonal Antibody; Anti-Cytokeratin 20/KRT20 Antibody; Rabbit Polyclonal; Cytokeratin 20/KRT20 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CD20 Antibody; Anti-CK-20 Antibody; Anti-CK20 Antibody; Anti-K20 Antibody; Anti-KRT21 Antibody; keratin 20; type I; anti-KRT20 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Cytokeratin 20/KRT20
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-KRT20 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human Cytokeratin 20/KRT20 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Cytokeratin 20/KRT20 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IF (Immunofluorescence)

Immunofluorescence staining of KRT20 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human KRT20 polyclonal antibody (dilution ratio 1:100) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human Cytokeratin 20 in human small intestine with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cytokeratin-20-krt20-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/257558>