

Rabbit anti-Human PDSS2 Polyclonal Antibody | anti-PDSS2 antibody

Anti-PDSS2 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257615	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PDSS2; Polyclonal Antibody; Anti-PDSS2 Antibody; Rabbit Polyclonal; PDSS2 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-bA59I9.3 Antibody; Anti-C6orf210 Antibody; Anti-COQ10D3 Antibody; Anti-DLP1 Antibody; Anti-hDLP1 Antibody; prenyl (solanesyl) diphosphate synthase; subunit 2; anti-PDSS2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human PDSS2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-PDSS2 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human PDSS2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human PDSS2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IP (Immunoprecipitation)

PDSS2 was immunoprecipitated using: Lane A: 0.5 mg HepG2 Whole Cell Lysate 4 uL anti-PDSS2 rabbit polyclonal antibody and 60 ug of Immnomagnetic beads Protein A/G. Primary antibody: Anti-PDSS2 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 44 kDa. Observed band size: 42 kDa.




IHC (Immunohistochemistry)

Immunochemical staining of human PDSS2 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human PDSS2 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-PDSS2 rabbit polyclonal antibody at 1:500 dilution Lane A: HepG2 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 44 kDa Observed band size: 42 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/pdss2-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/257615>